

Assessment Of Edentulism Among Outpatients- An Institutional Study

Neha Sharma M

Saveetha Dental College And Hospitals,
Saveetha Institute Of Medical And Technical Sciences,
Saveetha University,
Chennai- 77
E-Mail Id: 151701015.Sdc@Saveetha.Com

L. Leelavathi

Senior Lecturer
Department Of Public Health Dentistry
Saveetha Dental College And Hospitals,
Saveetha Institute Of Medical And Technical Science,
Saveetha University,
Chennai-77
E-Mail Id: Leelavathi.Sdc@Saveetha.Com

Abstract:

Background: The Total Rate Of Edentulism Is Said To Be Increasing In Developing Countries And This Has Been Attributed Mainly To The High Prevalence Of Periodontal Diseases And Caries. Several Reports Have Shown That The Prevalence Of Edentulism Is Higher In Males Than Females.

Aim: The Aim Of This Study Is To Assess The Edentulism Based On Acp Pdi Classification Among 65-74 Yrs Old Patients Visiting A Private Dental Hospital In Chennai

Materials And Methods: This Was A Comparative, Descriptive Study Done During April - June, 2020, Where All The Data Of The Patient Who Reported To The Dental Clinics In Saveetha Dental College, Simats, Chennai, India Was Obtained From The Department Of Public Health Dentistry. Patient Records Were Reviewed And Analysed Between June 2019 And March 2020. Data Was Collected And Tabulated. The Collected Data Was Further Analysed , Recorded In Microsoft Excel Software And Was Subjected To Chi Square Test For Statistical Analysis Using Ibm Spss Software.

Results: The Total Sample Size Of The Current Study Was 1,244 Cases. A Male Predilection Was Seen With Males Making Up 68% Of The Study Population. The Most Common Age Group That Was Observed Was 65-69 Years With 66%. Most Commonly Seen Acp Pdi Classification Was Class I With 70%. On Doing The Chi Square Test, It Was Found That Among The Males, Class I And Class Ii Edentulousness Was More Prevalent And Among The Females, Class Iii And Class Iv Edentulousness Was Found To Be More Prevalent. But It Was Found To Be Statistically Insignificant (P Value >0.05)

Conclusion: The Study Revealed That The Class I Acp Pdi Classification Was More Common Than Other Classes. There Was No Significant Association Between Edentulousness And Gender And Also Between Edentulousness And Age Among The Outpatients. Awareness Should Be Created Among The Public About The Factors Related To Tooth Loss And Also About The Treatment Options Availability To Treat The Edentulous State.

Keywords: Edentulousness; Gender; Acp Pdi ; Correlation; Age.

Introduction:

Edentulism is a debilitating and irreversible condition and is described as the “final marker of disease burden for oral health” [1]. Tooth loss can be identified by an edentulous space, which is a gap in the dental arch normally occupied by one tooth or more [2]. It could be partial or complete. A person may be partially edentulous or completely edentulous with all teeth missing in both upper and lower arch. One of the most common causes of tooth loss is dental caries [3] [4] [5] which commonly occur in pits and fissures of the occlusal surfaces in primary and permanent posterior teeth [6] [7]. Other causes for tooth loss are periodontal disease, physical injury or trauma [8], dietary habits [9] [10], and personal habits [11]. This can be prevented at an early stage by using sealants [12] and fluoridated products like fluoride varnish [13] [14] [15]. Although fluorides in excessive amounts can cause conditions like fluorosis [16]. Edentulousness has been found to increase in old age, which may partly be attributable to physical disabilities that could occur in old age [17]. The level of edentulousness is the main indicator for the oral health status of senior citizens. Some countries in Western Europe - Austria, France, Italy, Switzerland - had the lowest percentages of edentulous individuals, while the United Kingdom and the Netherlands had exceptionally high levels [13,18]. Data from several countries have consistently shown that edentulism is more prevalent among lower socioeconomic groups and among women, and that women have tended to become edentulous at a younger age [4,19]. Previous studies have also shown that several non-disease factors such as attitude, behavior, dental attendance, characteristics of the health care system and socio-demographic factors play important roles in the aetiopathogenesis of edentulism [20]. The ACP PDI (American College of Prosthodontists Prosthodontic Diagnostic Index) classification has been used as a diagnostic criteria in this present study for edentulousness. The edentulous span confined to a single arch and not exceeding 2 missing incisors in maxillary arch or 4 incisors in mandibular arch or 2 premolars, or 1 premolar and 1 molar in posterior region was categorized as Class I, single arch with canine missing or both arches with two teeth missing was categorized as Class II, one or two arches with three teeth missing or two molars missing was categorized as Class III and any maxillofacial defect was categorized as Class IV. [21]

Risk indicators of total tooth loss included age, gender, length of education, geographical area, urbanization, marital status, and medication for headache [22]. Previous studies also show that prevalence of edentulism was higher in the males than in the females in all age groups except in the 50-59 age group [23]. Another study showed that more number of men (52%) were partially edentulous when compared to women (48%), this could be because of the poor brushing habits due to their work schedule, and the need to seek treatment among the group of men were more when compared to women [24]. The level of edentulousness was found to be high, more so in rural than in urban people and more so in advancing age, with no significant difference between male and females [25]. Previously our team has a rich experience in working on various research projects across multiple disciplines [26-28][29-40].

A simple estimation of the edentulous condition is a rough indication of the prevalence of dental diseases and success or failure of dental care. Hence the aim of this study was to assess the edentulousness based on ACP PDI classification among 65-74 yr old patients visiting the private dental hospital.

Materials And Method:

This was an institutional study among patients with edentulousness who have visited Saveetha Dental College and hospitals in between June 2019- April 2020. The approval for this study setting was obtained from the institution ethics board and the study was conducted during April - June, 2020. The sample size of 1244 patients in which sampling bias was minimized with the verification of photos. The study was reviewed by two reviewers and it was cross verified. Records of patients presenting with edentulousness were reviewed and incomplete records in the system were excluded. ACP PDI classification was used to measure edentulousness which was one of the commonly used classification and hence assuring the internal validity of the study. ACP PDI classification system categorizes patients according to complexity of the edentulous condition and helps the clinician to make a diagnostically driven treatment plan. The case records of edentulous patients were collected by reviewing patient records and data of these patients was collected and tabulated. The following parameters including patients ID, age, gender, and ACP PDI classification were collected from the

Assessment Of Edentulism Among Outpatients- An Institutional Study

Records. Age Was Categorized Into Two Groups As 65-69 Years And 70-74 Years. After Further Verification By An External Reviewer, It Was Imported To Spss Software By Ibm For Statistical Analysis. Similar Methodology Was Employed In Other Cross Sectional Studies As Well [41]. Percentage , Mean, Standard Deviation, Frequency Of Parameters Were Analysed Using Descriptive Statistics.. Chi Square Test Was Used To Analyse Association Between Age And Gender With Edentulousness Classification. P Value Less Than 0.05 Was Considered To Be Statistically Significant.

Results:

In The Present Study, A Total Of 1,244 Patient Records Were Screened. Out Of Them, The Percentage Of Male Patients Was Found To Be 68% And Females Was Found To Be 32% [Figure 1]. Out Of The Total Sample Size, The Maximum Number Of Patients Belonged To The Age Group 65-69 Years (66%) And 34% Of The Patients Were Between 70-74 Years [Figure 2]. When Checked For The Acp Pdi Classification, 70% Of The Patients Showed Class I Edentulousness, 17.4% Of The Patients Showed Class Iii Edentulousness, About 10.5% Of The Patients Showed Class Iv Edentulousness And 1.2% Of Them Showed Class Ii [Figure 3]. On Correlating Edentulousness And Gender, Among The Males, 604 Patients Had Class I Edentulousness (71.5%), 12 Patients Had Class Ii Edentulousness (1.4%), 142 Patients Had Class Iii Edentulousness (16.8%), 87 Patients Had Class Iv Edentulousness (10.3%) And Among The Females, 278 Patients Had Class I Edentulousness (69.7%), 3 Patients Had Class Ii Edentulousness (0.8%), 74 Patients Had Class Iii Edentulousness (18.5%) And 44 Patients Had Class Iv Edentulousness (11%) [Figure 4]. On Doing The Chi Square Test, The P Value Was Found To Be $0.628 > 0.05$ Which Was Statistically Not Significant. On Correlating Age And Edentulousness, 589 Patients Between 65-69 Years Showed Class I Edentulousness (71.7%), 9 Patients Showed Class Ii (1.1%), 144 Patients Showed Class Iii Edentulousness (17.5%), 79 Patients Showed Class Iv Edentulousness (9.6%) And Among The Age Group 70-74 Years, 293 Patients Showed Class I Edentulousness (69.3%), 6 Patients Showed Class Ii Edentulousness (1.4%), 72 Patients Showed Class Iii Edentulousness (17%) And 52 Patients Showed Class Iv Edentulousness (12.3%) [Figure 5]. Although It Was Found To Be Statistically Not Significant. [P Value $0.491 > 0.05$].

Discussion:

The Complex Interaction Between Dental Diseases, The Tendency To Use Dental Care, Dental Attitude, And Affordability Of Non-Extraction Treatment Have Been Related To The Incidence Of Tooth Loss [42]. Oral Health Is Related To Diet In Many Ways, For Example, Nutritional Influences On Craniofacial Development, Oral Cancer And Oral Infectious Diseases[43] [44]; Poor Education, A Risk Factor For Poverty, Has Been Identified As A Major Factor In Edentulism . The Observed Prevalence Of Edentulism Was Moderate To Low In The Adult And Elderly Groups [45]. Clinically, Partial Edentulism Results In Drifting And Tilting Of Adjacent Teeth , Supra Eruption Of Opposing Teeth, Altered Speech, Changes In Facial Appearance And Tempero-Mandibular Disorders [46]. Edentulism Has A Significant Impact On Health And The Overall Quality Of Life. Studies On Self-Perception Have Demonstrated That Tooth Loss Is Associated With Aesthetical, Functional, Psychological And Social Impacts On Individuals. Moreover; There Is Sufficient Evidence To Indicate That Loss Of Teeth Can Adversely Affect Food Selection And Ultimately Result In Incidence Of Various Health Disorders [47].

Previous Literature Shows That Women Were More Likely To Be Edentulous [48]. Another Study Shows That Females And Unemployed Groups Have More Edentulousness And Upper Arch Was More Common [49]. Whereas Another Study Shows That There Is More Prevalence Of Edentulousness In Males 64% Than In Females 35% [50]. This Is Because Females Take More Care Of Their Oral Health Than Males And In Rural Areas [51]. A Previous Study Shows That The Women Show A Higher Proportion Of Edentulousness [52]. Edentulism Was Not Associated With Total Energy Or Food Intake But Was Associated With The Food Groups Consumed , Particularly Fat, Micronutrients, And Hard-To-Chew Foods [53]. Another Study Shows That Edentulism Is More Common Among Those With Less Education And Income [54]. Our Institution Is Passionate About High Quality Evidence Based Research And Has Excelled In Various Fields ([55-65].

Awareness About Medical And Dental Negligence Among The Public Is Growing In India [66]. Hence Dental Education Should Be Targeted At The Uneducated Populace, The Rural Dwellers And Low-Income Groups To Reduce The Rate Of Total Edentulism.

Conclusion:

Class I Acp Pdi Classification Was More Common Among The Study Population. Class I And Class Ii Edentulousness Based On Acp Pdi Classification Was Found To Be More Prevalent In Males Than Females Whereas Class Iii And Class Iv Edentulousness Were More Prevalent Among Females. Class I And Class Iii Edentulousness Were More Prevalent Among The Age Group 65-69 Years, Whereas Class Ii And Class Iv Edentulousness Were More Prevalent Among The Age Group 70-74 Years, But These Were Not Statistically Significant.

Acknowledgement:

The Authors Would Like To Acknowledge The Help And Support Rendered By Information Technology Department Of Saveetha Dental College And Hospitals And The Management For Their Constant Assistance With The Research.

Author Contributions:

Author 1 (Neha Sharma M) Carried Out The Retrospective Study By Collecting Data And Drafted The Manuscript After Performing The Necessary Statistical Analysis. Author 2 (Dr. L. Leelavathi) Aided In Conception Of The Topic, Has Participated In The Study Design, Statistical Analysis And Has Supervised In The Preparation Of The Manuscript. Author 3 (Dr. L. Leelavathi) Has Participated In The Study Design, And Has Coordinated In Developing The Manuscript. All The Authors Have Discussed The Results Among Themselves And Contributed To The Final Manuscript.

Conflict Of Interest:

Nil

References:

- [1] Emami E, De Souza Rf, Kabawat M, Feine Js. The Impact Of Edentulism On Oral And General Health. *International Journal Of Dentistry* 2013;2013:1–7. <https://doi.org/10.1155/2013/498305>.
- [2] Madhankumar S, Mohamed K, Natarajan S, Kumar V, Athiban I, Padmanabhan Tv. Prevalence Of Partial Edentulousness Among The Patients Reporting To The Department Of Prosthodontics Sri Ramachandra University Chennai, India: An Epidemiological Study. *Journal Of Pharmacy And Bioallied Sciences* 2015;7:643. <https://doi.org/10.4103/0975-7406.163580>.
- [3] Prabakar J, John J, Srisakthi D. Prevalence Of Dental Caries And Treatment Needs Among School Going

Assessment Of Edentulism Among Outpatients- An Institutional Study

- Children Of Chandigarh. *Indian J Dent Res* 2016;27:547–52.
- [4] Mathew Mg, Samuel Sr, Soni Aj, Roopa Kb. Evaluation Of Adhesion Of Streptococcus Mutans, Plaque Accumulation On Zirconia And Stainless Steel Crowns, And Surrounding Gingival Inflammation In Primary Molars: Randomized Controlled Trial. *Clinical Oral Investigations* 2020. <https://doi.org/10.1007/S00784-020-03204-9>.
- [5] Samuel Sr, Acharya S, Rao Jc. School Interventions–Based Prevention Of Early-Childhood Caries Among 3–5-Year-Old Children From Very Low Socioeconomic Status: Two-Year Randomized Trial. *Journal Of Public Health Dentistry* 2020;80:51–60. <https://doi.org/10.1111/Jphd.12348>.
- [6] Mohapatra S, Pradeep Kumar R, Meignana Arumugham I, Sri Sakthi D, Jayashri P. Assessment Of Microhardness Of Enamel Carious Like Lesions After Treatment With Nova Min, Bio Min And Remin Pro Containing Toothpastes: An In Vitro Study. *Indian Journal Of Public Health Research & Development* 2019;10:375. <https://doi.org/10.5958/0976-5506.2019.02832.8>.
- [7] Prabhakar Ar, Murthy S, Sugandhan S. Comparative Evaluation Of The Length Of Resin Tags, Viscosity And Microleakage Of Pit And Fissure Sealants - An In Vitro Scanning Electron Microscope Study. *Contemporary Clinical Dentistry* 2011;2:324. <https://doi.org/10.4103/0976-237x.91797>.
- [8] Leelavathi L, Karthick R, Sankari S, Babu N. Avulsed Tooth – A Review. *Biomedical And Pharmacology Journal* 2016;9:847–50. <https://doi.org/10.13005/Bpj/1015>.
- [9] Neralla M, Jayabalan J, George R, Rajan J, P Skm, Haque Ae, Et Al. Role Of Nutrition In Rehabilitation Of Patients Following Surgery For Oral Squamous Cell Carcinoma. *International Journal Of Research In Pharmaceutical Sciences* 2019;10:3197–203. <https://doi.org/10.26452/Ijrps.V10i4.1622>.
- [10] Pratha Aa, Ashwatha Pratha A, Prabakar J. Comparing The Effect Of Carbonated And Energy Drinks On Salivary Ph- In Vivo Randomized Controlled Trial. *Research Journal Of Pharmacy And Technology* 2019;12:4699. <https://doi.org/10.5958/0974-360x.2019.00809.6>.
- [11] Harini G, Leelavathi L. Nicotine Replacement Therapy For Smoking Cessation–An Overview. *Indian Journal Of Public Health Research & Development* 2019;10:3588. <https://doi.org/10.5958/0976-5506.2019.04144.5>.
- [12] Prabakar J, John J, Arumugham I, Kumar R, Srisakthi D. Comparative Evaluation Of Retention, Cariostatic Effect And Discoloration Of Conventional And Hydrophilic Sealants - A Single Blinded Randomized Split Mouth Clinical Trial. *Contemporary Clinical Dentistry* 2018;9:233. https://doi.org/10.4103/Ccd.Ccd_132_18.
- [13] Kumar Rp, Pradeep Kumar R, Vijayalakshmi B. Assessment Of Fluoride Concentration In Ground Water In Madurai District, Tamil Nadu, India. *Research Journal Of Pharmacy And Technology* 2017;10:309. <https://doi.org/10.5958/0974-360x.2017.00063.4>.
- [14] Khatri S, Madan K, Srinivasan S, Acharya S. Retention Of Moisture-Tolerant Fluoride-Releasing Sealant And Amorphous Calcium Phosphate-Containing Sealant In 6–9-Year-Old Children: A Randomized Controlled Trial. *Journal Of Indian Society Of Pedodontics And Preventive Dentistry* 2019;37:92. https://doi.org/10.4103/Jisppd.Jisppd_173_18.
- [15] Prabakar J, John J, Arumugham I, Kumar R, Sakthi D. Comparing The Effectiveness Of Probiotic, Green Tea, And Chlorhexidine- And Fluoride-Containing Dentifrices On Oral Microbial Flora: A Double-Blind, Randomized Clinical Trial. *Contemporary Clinical Dentistry* 2018;9:560. https://doi.org/10.4103/Ccd.Ccd_659_18.
- [16] Kumar Rp, Pradeep Kumar R, Preethi R. Assessment Of Water Quality And Pollution Of Porur, Chembarambakkam And Puzhal Lake. *Research Journal Of Pharmacy And Technology* 2017;10:2157. <https://doi.org/10.5958/0974-360x.2017.00380.8>.
- [17] Fouda Sm, Al-Harbi Fa, Khan Sq, Virtanen Ji, Raustia A. Missing Teeth And Prosthetic Treatment In Patients Treated At College Of Dentistry, University Of Dammam. *Int J Dent* 2017;2017:7593540.
- [18] Bourgeois D, Nihtila A, Mersel A. Prevalence Of Caries And Edentulousness Among 65-74-Year-Olds In Europe. *Bull World Health Organ* 1998;76:413–7.
- [19] Eklund Sa, Burt Ba. Risk Factors For Total Tooth Loss In The United States; Longitudinal Analysis Of National Data. *Journal Of Public Health Dentistry* 1994;54:5–14. <https://doi.org/10.1111/J.1752-7325.1994.Tb01173.X>.
- [20] Esan Ta, Olusile Ao, Akeredolu Pa, Esan Ao. Socio-Demographic Factors And Edentulism: The Nigerian Experience. *Bmc Oral Health* 2004;4. <https://doi.org/10.1186/1472-6831-4-3>.
- [21] Sharma A. Description Of Partial Edentulism Among Patients Visiting The Prosthodontic N.D.
- [22] Suominen-Taipale A-L, Alanen P, Helenius H, Nordblad A, Uutela A. Edentulism Among Finnish Adults Of Working Age, 1978?1997. *Community Dentistry And Oral Epidemiology* 1999;27:353–65. <https://doi.org/10.1111/J.1600-0528.1999.Tb02032.X>.
- [23] Hoover Jn, Mcdermott Re. Edentulousness In Patients Attending A University Dental Clinic. *J Can Dent Assoc*

- 1989;55:139–40.
- [24] Kumar C, Manimaran P, Saisadan D, Abirami M, Kumar N, Mani J. Partial Edentulousness In A Rural Population Based On Kennedy's Classification: Epidemiological Study. *Journal Of Pharmacy And Bioallied Sciences* 2017;9:34. https://doi.org/10.4103/jpbs.jpbs_87_17.
- [25] Shah N, Parkash H, Sunderam Kr. Edentulousness, Denture Wear And Denture Needs Of Indian Elderly--A Community-Based Study. *J Oral Rehabil* 2004;31:467–76.
- [26] Hafeez N, Others. Accessory Foramen In The Middle Cranial Fossa. *Research Journal Of Pharmacy And Technology* 2016;9:1880.
- [27] Krishnan Rp, Ramani P, Sherlin Hj, Sukumaran G, Ramasubramanian A, Jayaraj G, Et Al. Surgical Specimen Handover From Operation Theater To Laboratory: A Survey. *Ann Maxillofac Surg* 2018;8:234–8.
- [28] Somasundaram S, Ravi K, Rajapandian K, Gurunathan D. Fluoride Content Of Bottled Drinking Water In Chennai, Tamilnadu. *J Clin Diagn Res* 2015;9:Zc32.
- [29] Felicita As, Sumathi Felicita A. Orthodontic Extrusion Of Ellis Class Viii Fracture Of Maxillary Lateral Incisor – The Sling Shot Method. *The Saudi Dental Journal* 2018;30:265–9. <https://doi.org/10.1016/j.sdentj.2018.05.001>.
- [30] Kumar S, Rahman R. Knowledge, Awareness, And Practices Regarding Biomedical Waste Management Among Undergraduate Dental Students. *Asian J Pharm Clin Res* 2017;10:341.
- [31] Gurunathan D, Shanmugaavel Ak. Dental Neglect Among Children In Chennai. *J Indian Soc Pedod Prev Dent* 2016;34:364.
- [32] Sneha S, Others. Knowledge And Awareness Regarding Antibiotic Prophylaxis For Infective Endocarditis Among Undergraduate Dental Students. *Asian Journal Of Pharmaceutical And Clinical Research* 2016:154–9.
- [33] Dhinesh B, Isaac Joshuaramesh Lalvani J, Parthasarathy M, Annamalai K. An Assessment On Performance, Emission And Combustion Characteristics Of Single Cylinder Diesel Engine Powered By Cymbopogon Flexuosus Biofuel. *Energy Convers Manage* 2016;117:466–74.
- [34] Choudhari S, Thenmozhi Ms. Occurrence And Importance Of Posterior Condylar Foramen. *Laterality* 2016;8:11–43.
- [35] Paramasivam A, Vijayashree Priyadharsini J, Raghunandhakumar S. N6-Adenosine Methylation (M6a): A Promising New Molecular Target In Hypertension And Cardiovascular Diseases. *Hypertens Res* 2020;43:153–4.
- [36] Wu F, Zhu J, Li G, Wang J, Veeraraghavan Vp, Krishna Mohan S, Et Al. Biologically Synthesized Green Gold Nanoparticles From Siberian Ginseng Induce Growth-Inhibitory Effect On Melanoma Cells (B16). *Artif Cells Nanomed Biotechnol* 2019;47:3297–305.
- [37] Palati S, Ramani P, Shrelin H, Sukumaran G, Ramasubramanian A, Don Kr, Et Al. Knowledge, Attitude And Practice Survey On The Perspective Of Oral Lesions And Dental Health In Geriatric Patients Residing In Old Age Homes. *Indian Journal Of Dental Research* 2020;31:22. https://doi.org/10.4103/ijdr.ijdr_195_18.
- [38] Saravanan M, Arokiyaraj S, Lakshmi T, Pugazhendhi A. Synthesis Of Silver Nanoparticles From Phenerochaete Chrysosporium (Mtcc-787) And Their Antibacterial Activity Against Human Pathogenic Bacteria. *Microb Pathog* 2018;117:68–72.
- [39] Govindaraju L, Gurunathan D. Effectiveness Of Chewable Tooth Brush In Children-A Prospective Clinical Study. *J Clin Diagn Res* 2017;11:Zc31.
- [40] Vijayakumar Jain S, Muthusekhar Mr, Baig Mf, Senthilnathan P, Loganathan S, Abdul Wahab Pu, Et Al. Evaluation Of Three-Dimensional Changes In Pharyngeal Airway Following Isolated Lefort One Osteotomy For The Correction Of Vertical Maxillary Excess: A Prospective Study. *J Maxillofac Oral Surg* 2019;18:139–46.
- [41] Kannan Ssd, Kumar Vs, Rathinavelu Pk, Indiran Ma. Awareness And Attitude Towards Mass Disaster And Its Management Among House Surgeons In A Dental College And Hospital In Chennai, India. *Disaster Management And Human Health Risk V* 2017. <https://doi.org/10.2495/Dman170121>.
- [42] Vadavadagi Sy, Srinivasa H, Goutham Gb, Hajira N, Lahari M, Reddy Gtp. Partial Edentulism And Its Association With Socio-Demographic Variables Among Subjects Attending Dental Teaching Institutions, India. *J Int Oral Health* 2015;7:60–3.
- [43] L L, Leelavathi L, Thoudam S, Anitha M. Nutrition And Oral Health - (Review). *Biomedical And Pharmacology Journal* 2015;8:545–8. <https://doi.org/10.13005/bpj/744>.
- [44] Pavithra Rp, Preethi Pavithra R, Jayashri P. Influence Of Naturally Occurring Phytochemicals On Oral Health. *Research Journal Of Pharmacy And Technology* 2019;12:3979. <https://doi.org/10.5958/0974-360x.2019.00685.1>.
- [45] Medina-Solís Ce, Pérez-Núñez R, Maupomé G, Casanova-Rosado Jf. Edentulism Among Mexican Adults Aged

Assessment Of Edentulism Among Outpatients- An Institutional Study

- 35 Years And Older And Associated Factors. *American Journal Of Public Health* 2006;96:1578–81. <https://doi.org/10.2105/Ajph.2005.071209>.
- [46] Jeyapalan V. Partial Edentulism And Its Correlation To Age, Gender, Socio-Economic Status And Incidence Of Various Kennedy's Classes– A Literature Review. *Journal Of Clinical And Diagnostic Research* 2015. <https://doi.org/10.7860/Jcdr/2015/13776.6124>.
- [47] D'Souza K, Aras M. Association Between Socio-Demographic Variables And Partial Edentulism In The Goan Population: An Epidemiological Study In India. *Indian Journal Of Dental Research* 2014;25:434. <https://doi.org/10.4103/0970-9290.142519>.
- [48] Shamdol Z, Ismail Nm, Hamzah Nt, Ismail Ar. Prevalence And Associated Factors Of Edentulism Among Elderly Muslims In Kota Bharu, Kelantan, Malaysia. *Journal Of The Islamic Medical Association Of North America* 2008;40. <https://doi.org/10.5915/40-4-4431>.
- [49] Sapkota B, Adhikari B, Upadhya C. A Study Of Assessment Of Partial Edentulous Patients Based On Kennedy's Classification At Dhulikhel Hospital Kathmandu University Hospital. *Kathmandu University Medical Journal* 2015;11:325–7. <https://doi.org/10.3126/Kumj.V11i4.12542>.
- [50] Patturaja K, Leelavathi L, Jayalakshmi S. Choice Of Rotary Instrument Usage Among Endodontists – A Questionnaire Study. *Biomedical And Pharmacology Journal* 2018;11:851–6. <https://doi.org/10.13005/Bpj/1441>.
- [51] Arandi N, Mustafa S. The Prevalence Of Different Kennedy's Classifications And Their Relationship With Age, Gender, And Location Of The Edentulous Arches In Palestine. *International Journal Of Preventive And Clinical Dental Research* 2019;6:53. https://doi.org/10.4103/Inpc.Inpc_51_19.
- [52] Patel Jy. Assessment Of Partially Edentulous Patients Based On Kennedy's Classification And Its Relation With Gender Predilection N.D.
- [53] Caplan Dj, Weintraub Ja. The Oral Health Burden In The United States: A Summary Of Recent Epidemiologic Studies. *J Dent Educ* 1993;57:853–62.
- [54] Lee Js, Weyant Rj, Corby P, Kritchevsky Sb, Harris Tb, Rooks R, Et Al. Edentulism And Nutritional Status In A Biracial Sample Of Well-Functioning, Community-Dwelling Elderly: The Health, Aging, And Body Composition Study. *The American Journal Of Clinical Nutrition* 2004;79:295–302. <https://doi.org/10.1093/Ajcn/79.2.295>.
- [55] Vijayashree Priyadharsini J. In Silico Validation Of The Non-Antibiotic Drugs Acetaminophen And Ibuprofen As Antibacterial Agents Against Red Complex Pathogens. *J Periodontol* 2019;90:1441–8.
- [56] Pc J, Marimuthu T, Devadoss P. Prevalence And Measurement Of Anterior Loop Of The Mandibular Canal Using Cbct: A Cross Sectional Study. *Clin Implant Dent Relat Res* 2018.
- [57] Ramesh A, Varghese S, Jayakumar Nd, Malaiappan S. Comparative Estimation Of Sulfiredoxin Levels Between Chronic Periodontitis And Healthy Patients - A Case-Control Study. *J Periodontol* 2018;89:1241–8.
- [58] Ramadurai N, Gurunathan D, Samuel Av, Subramanian E, Rodrigues Sjl. Effectiveness Of 2% Articaine As An Anesthetic Agent In Children: Randomized Controlled Trial. *Clin Oral Investig* 2019;23:3543–50.
- [59] Sridharan G, Ramani P, Patankar S, Vijayaraghavan R. Evaluation Of Salivary Metabolomics In Oral Leukoplakia And Oral Squamous Cell Carcinoma. *J Oral Pathol Med* 2019;48:299–306.
- [60] Ezhilarasan D, Apoorva Vs, Ashok Vardhan N. Syzygium Cumini Extract Induced Reactive Oxygen Species-Mediated Apoptosis In Human Oral Squamous Carcinoma Cells. *J Oral Pathol Med* 2019;48:115–21.
- [61] Mathew Mg, Samuel Sr, Soni Aj, Roopa Kb. Evaluation Of Adhesion Of Streptococcus Mutans, Plaque Accumulation On Zirconia And Stainless Steel Crowns, And Surrounding Gingival Inflammation In Primary Molars: Randomized Controlled Trial. *Clin Oral Investig* 2020:1–6.
- [62] Samuel Sr. Can 5-Year-Olds Sensibly Self-Report The Impact Of Developmental Enamel Defects On Their Quality Of Life? *Int J Paediatr Dent* 2021;31:285–6.
- [63] R H, Hannah R, Ramani P, Ramanathan A, R Jm, Gheena S, Et Al. Cyp2 C9 Polymorphism Among Patients With Oral Squamous Cell Carcinoma And Its Role In Altering The Metabolism Of Benzo[A]Pyrene. *Oral Surgery, Oral Medicine, Oral Pathology And Oral Radiology* 2020;130:306–12. <https://doi.org/10.1016/J.Ooooo.2020.06.021>.
- [64] Chandrasekar R, Chandrasekhar S, Sundari Kks, Ravi P. Development And Validation Of A Formula For Objective Assessment Of Cervical Vertebral Bone Age. *Prog Orthod* 2020;21:38.
- [65] Vijayashree Priyadharsini J, Smiline Girija As, Paramasivam A. In Silico Analysis Of Virulence Genes In An Emerging Dental Pathogen A. *Baumannii* And Related Species. *Arch Oral Biol* 2018;94:93–8.
- [66] Srudhy L, Anitha M. Medicolegal Case Scenarios In Dental Practice - A Review Of Literature. *Biomedical And Pharmacology Journal* 2015;8:537–41. <https://doi.org/10.13005/Bpj/742>.

Figure Titles:

Fig 1: Bar Graph Representing The Frequency Of Occurrence Of Edentulousness Among Different Genders

Fig 2: Bar Graph Representing The Frequency Of Occurrence Of Edentulousness Among Different Age Groups

Fig 3: Bar Graph Representing The Acp Pdi Classification Of Edentulous Patients

Fig 4: Bar Graph Representing Association Between Gender And Edentulousness Among Patients

Fig 5: Bar Graph Representing Association Between Age And Edentulousness Among Patients

Figures:

Assessment Of Edentulism Among Outpatients- An Institutional Study




Figure 1: Bar Graph Representing The Frequency Of Occurrence Of Edentulousness Among Different Genders. Y Axis Represents The Count Of Patients With Edentulousness And X Axis Represents The Gender Of The Patients. The Colour Purple Denotes Male Patients And Yellow Denotes Female Patients. The Above Graph Shows That 68% Of The Patients Were Males And 32% Were Females Suggesting Males Were The More Prevalent Study Participants.




Figure 2: Bar Graph Representing The Frequency Of Occurrence Of Edentulousness Among Different Age Groups. Y Axis Represents The Count Of Patients With Edentulousness And X Axis Represents The Age Group Of The Patients. The Colour Blue Denotes The Age Group 65-69 Years And Green Colour Denotes Patients Between 70-74 Years. The Above Graph Shows That 66% Of The Patients Are Between 65-69 Years And 34% Of The Patients Are Between 70-74 Years. Thus The Age Group 65-69 Years Was Found To Be More Prevalent.

Assessment Of Edentulism Among Outpatients- An Institutional Study




Figure 3: Bar Graph Representing The Acp Pdi Classification Of Edentulous Patients. Y Axis Represents The Count Of Patients With Edentulousness And X Axis Represents The Acp Pdi Classification Of Edentulous Patients. The Colour Grey Denotes Class I, Light Blue Colour Denotes Class Ii, Light Green Denotes Class Iii And Brown Colour Denotes Class Iv Edentulousness. The Above Graph Shows That 71% Of The Patients Have Class I Edentulousness, 1.2% Of The Patients Have Class Ii Edentulousness ,17.3% Of The Patients Have Class Iii Edentulousness, And 10.5% Of The Patients Have Class Iv Edentulousness. Class I Edentulousness Was More Prevalent Among The Study Population.




Figure 4: Bar Graph Representing Association Between Gender And Edentulousness Among Patients. X Axis Represents The Acp Pdi Classification Of Edentulousness And Y Axis Represents The Gender Of The Edentulous Patients. The Colour Purple Denotes Males And Yellow Denotes Females Patients. The Above Graph Shows That Class I [604(71.5%)] And Classii Edentulousness [12 (1.4%)] Were More Prevalent Among Males Whereas Class Iii [74(18.5%)] And Class Iv Edentulousness [44(11%)] Were More Prevalent Among Females. Chi Square Test Showed There Was No Significant Association Between Gender And Edentulousness Among 65-74 Yr Old Patients Visiting The Private Dental Hospital ; Pearson Chi Square Value- 1.740, P Value- 0.628 (>0.05). Classii Edentulousness [12 (1.4%)] Were More Prevalent Among Males When Compared To Other Classes.

Assessment Of Edentulism Among Outpatients- An Institutional Study




Figure 5: Bar Graph Representing Correlation Between Age And Edentulousness Among Patients. X Axis Represents The Acp Pdi Classification Of Edentulousness And Y Axis Represents The Age Group Of The Edentulous Patients. The Colour Blue Denotes The Age Group 65-69 Years And Green Colour Denotes Patients Between 70-74 Years. The Above Graph Shows That Class I And Class Iii Edentulousness Were More Prevalent Among The Age Group 65-69 Years [589 (71.7%) And 144(17.5%)] Respectively, Class Ii And Class Iv Edentulousness Were More Prevalent Among The Age Group 70-74 Years [6 (1.4%) And 52(12.3%)] Respectively. Chi Square Test Showed That There Was No Significant Association Between Age And Edentulousness Among 65-74 Yr Old Patients Visiting The Private Dental Hospital ; Pearson Chi Square Value- 2.416, P Value- 0.491(>0.05).