

Prevalence And Analysis Of Factors Associated With Canine Impaction In Class Iii Malocclusion. - A Retrospective Case Study

Amina Mehrin Bano

Saveetha Dental College
Saveetha Institute Of Medical And Technical Sciences
Saveetha University
Chennai
Email: 151501085.Sdc@Saveetha.Com

Srirengalakshmi

Senior Lecturer ,
Department Of Orthodontics ,
Saveetha Dental College And Hospital
Saveetha Institute Of Medical And Technical Sciences
Saveetha University
Chennai
Email: Srirengalakshmi.Sdc@Saveetha.Com

Abstract :

Impacted Teeth Are Those With A Delayed Eruption Time Or That Are Not Expected To Erupt Completely Based On Clinical And Radiographic Assessment. After The Third Molars, The Maxillary Canine Is The Second Most Frequently Impacted Tooth In The Dental Arch. The Prevalence Of Impacted Maxillary Canine Is 0.9 To 2.2%. Knowledge Of Occlusion Of Each Patient Can Contribute Significantly To Complete Care And Instruction. The Aim Of The Study Was To Analyse The Association Of Canine Impaction To Age , Gender And Types In Class Iii Malocclusions. This Was A Single Retrospective Study Set In A University Dental Hospital With Predominantly South Indian Population. The Data Was Collected From The Electronic Database Of The University. A Total Of 50 Patients Were Included In The Study .Data Tabulation Was Done In Excel Then It Was Imported And Assessed Using Statistical Package For Social Science 20(Spss,Ibm Corporation). A Test For Frequency Of Incidence Of Impacted Canine Among Patients Diagnosed With Class Iii Malocclusions Was Conducted. Chi-Square Test Was Conducted For Association Of Age , Gender , Type Of Class Iii Malocclusion. The Results Were Descriptively Presented In The Form Of Graphs And Tables . Only Patients With True Class Iii Malocclusion In The Age Range Of 13-30 Yrs Were Included In The Study. Among The Study Population 78% Of The Patients Were Diagnosed With A Skeletal Class Iii And 22% Of The Patients Were Diagnosed With A Dental Class Iii. Out Of The Whole Study Population Only Four Patients Had Impacted Canines That Is 8% Of The Total Study Population. Of The Total Study Population , 6% Of The Patients With Impacted Canines Were Male And 2% Of The Patients Were Female. Canine Impaction Has Statistically Significant Association To Both Adult And Child Age Groups, $P=0.028$. Also Males Patients With Class Iii Malocclusions Showed A Higher Association To Canine Impaction , $P=0.004$. Among The Study Population , Canine Impaction Showed More Association With Dental Class Iii Patients , ($P = 0.038$).

Keywords : Canine Impaction ; Class Iii Malocclusion ; Incidence ; Skeletal Class Iii ; Dental Class Iii

Introduction

Impacted Teeth Are Those With A Delayed Eruption Time Or That Are Not Expected To Erupt Completely Based On Clinical And Radiographic Assessment. [1] Failure Of Tooth Eruption May Be The Consequence Of Local Factors. These Factors May Include Mechanical Obstruction; Insufficient Space In The Dental Arch Due To Skeletal Discrepancy (Micrognathia); Or To The Premature Loss Of Deciduous Teeth Or To Tooth Arch Size Discrepancy. Systemic Factors Such As Genetic Disorders, Endocrine Deficiencies, Previous Irradiation Of The Jaw May Also Be

Prevalence And Analysis Of Factors Associated With Canine Impaction In Class Iii Malocclusion. - A Retrospective Case Study

Associated With The Failure Of Tooth Eruption . In A Majority Of Systemic Conditions Multiple Teeth Are Usually Impacted. In Most Cases However The Specific Cause Of Failure Of Eruption Remains Unknown.[2]

Third Molars, Maxillary Canine, Maxillary And Mandibular Premolar, And Maxillary Central Incisor Are The Teeth Most Frequently Involved. After The Third Molars, The Maxillary Canine Is The Second Most Frequently Impacted Tooth In The Dental Arch. The Prevalence Of Impacted Maxillary Canine Is 0.9 To 2.2% But Mandibular Canine Impact Impaction Occurs Less Frequently.[3,4] A Large Number Of Completely Impacted Teeth May Be Retained When Asymptomatic. However, Bishara Et Al, Suggested A Few Sequelae Of Canine Impaction, Such As Labial Or Lingual Malpositioning Of The Impacted Tooth, Migration Of The Neighbouring Teeth And Loss Of Arch Length, External Root Resorption Of The Impacted Tooth As Well As The Neighbouring Tooth, Infection Particularly With Partial Eruption Resulting In Pain And Trismus And Referred Pain. [5]

Occlusion Is The Relationship Between The Maxillary And Mandibular Teeth When They Approach Each Other, Either During Chewing Or At Rest.[6] It Is Examined And Recorded As Part Of The Oral Examination. Knowledge Of Occlusion Of Each Patient Can Contribute Significantly To Complete Care And Instruction. Recognition Of Malocclusion Assists In The Reference Of Patients To The Orthodontist, Gives Many Valuable Points Of Reference For Patient Instruction, And Determines Necessary Adaptations In Technique.[7] Occlusion Has Been Defined As The Interdigitation Of Maxillary And Mandibular Teeth [8]. This Description Is Misleading In That It Implies That Occlusion Is A Static Contact Relationship Or Arrangement Of Teeth. The Practical Concept Of Occlusion Is To Recognise The Interplay Between The Teeth, Temporomandibular Joint And Neuromuscular System.[9] The World Health Organisation (1962) Define Malocclusion As A Dentofacial Anomaly.

Currently , Even Though Malocclusions Occurs In A Majority Of The Population , It's A Variation To Normal Occlusion And Is An Eccentric Condition That Required Treatment . [8] According To The Angle Classification Class Iii Malocclusion Is Defined As The Mesially Placed Lower Molar In Relation To The Upper Molar With No Specifics In Regard To The Line Of Occlusion [10,11] . Nevertheless As With All Angles Classification, Class Iii Malocclusion Comprises Several Skeletal And Dental Components That May Differ From The Concept Of Normality. Previous Studies Regarding Force Implementation , Resin Penetration , Soft Tissue Profiles , Mini Implants For Anchorage [12–22] , Clinical Reports, Clinical Trials Regarding Different Approaches [23–26] Kindled Our Interest In Epidemiological Studies Related To Malocclusion And Impactions . Multiple Studies Have Documented The Prevalence Of Class Iii Malocclusion. Although The Incidence Of Canine Impaction And Its Relation To Class Iii Malocclusion Has Not Been Investigated In The South Indian Population. Previously Our Team Has A Rich Experience In Working On Various Research Projects Across Multiple Disciplines The [27–29][30–41] The Aim Was To Analyse The Association Of Canine Impaction To Age , Gender And Types In Class Iii Malocclusions .

Materials And Methods :

This Was A Retrospective Study Conducted In A University Dental Hospital With Predominantly South Indian Population. The Data Was Collected From The Electronic Database Of The University. The Ease Of Access And The Availability Of A Standardised Data Collection Format, Complete Availability Of Photographic Data Of Both Pre And Post Orthodontic Therapy Was An Advantage. The Approval For The Research Was Given By The Ethical Review Board Of Saveetha Institute Of Medical And Technical Sciences (Sdc/Sihc/2020/Diasdata/0619-032020).

The Study Population Selected Were Outpatients Who Were Diagnosed With A Class Iii Malocclusion At The Department Of Orthodontics At Saveetha Dental College From The Duration Of June 2019 To March 2020. Patients With Trauma To Teeth , Systemic Diseases, Previous Orthodontic Treatment, Periodontal Pathology Were Excluded From The Study. All Incomplete Data Was Excluded On The Basis Of Possible Bias. As The Data Was Collected From The University Database, And As It Was Verified Both By Department And The Patient And Cross-Checked With The Photographic Record, The Study Was Said To Be Internally Valid. The Results Of This Study Can Be Applied To The South Indian Population For Epidemiological Inferences.

A Total Of 50 Patients Were Included In The Study. The Data Tabulated From The University Database Was Pid (Patient Identification), Duration Of Treatment , Age, Gender, Pret-Op Digital Records Etc. Data Tabulation Was Done In Excel. The Data Was Imported And Transcribed In Statistical Package For Social Science 20 (Spss, Ibm Corporation). A Test For Frequency Of Incidence Of Impacted Canine Was Conducted. Chi-Square Test Was Conducted For Association Of Age , Gender , Type Of Class Iii Malocclusion . Finally, The Results Were Descriptively Presented.

Results :

A Total Number Of 50 Patients Were Included In The Study Who Were Diagnosed With Class Iii Malocclusion During The Duration Of The Study. The Patients Included Aged From A Range Of 12 Years To 31 Years . Only Patients With True Class Iii Malocclusion Were Included In The Study. Of The Whole Study Population , 78% Of The Patients Were Diagnosed With A Skeletal Class Iii And 22% Of The Patients Were Diagnosed With A Class Iii Subdivision. Out Of The Whole Study Population Only Four Patients Had Impacted Canines That Is 8% Of The Total Study Population. 6% Of The Patients With Impacted Canine Were Male And 2% Of The Patients Were Female. Canine Impaction Has Statistically Significant Association Among Both Adult And Child Age Groups, $P=0.028$ (Graph 3) Also Males Patients With Class Iii Malocclusions Showed A Higher Association To Canine Impaction , $P=0.004$ (Graph 1) . Among The Study Population , Canine Impaction Showed More Association With Dental Class Iii Patients , $P = 0.038$.(Graph 2)

Discussion :

In Our Retrospective Study , 50 Patients Who Had Been Diagnosed With Class Iii Malocclusions Had Been Radiographically Evaluated For Presence Of Impacted Canines . Among The 50 Patients Diagnosed With Class Iii Malocclusion , Only 4 Patients , That Is 8% Of The Population Had Impacted Canines . Out Of Which Male Patients Showed A Higher Prevalence 6% And 2% Of The Female Patients Had Impacted Canines(Graph 1) . Canine Impaction Has Statistically Significant Association Among Both Adult And Child Age Groups, $P=0.028$ (Graph 3) Also Males Patients With Class Iii Malocclusions Showed A Higher Association To Canine Impaction , $P=0.004$ (Graph 1) . Among The Study Population , Canine Impaction Showed More Association With Dental Class Iii Patients , $P = 0.038$.(Graph 2). The Large Variation In Angle Class Iii Malocclusion Prevalence Rates Can Be Seen Globally Suggesting A High Level Of Variability Across Geographic Regions And Races. Chinese And Malaysian Groups Showed A Much Higher Mean Prevalence Rate Than Other Racial Groups: 15.69% And 16.59%, Respectively. This Is Also Consistent With Previous Reports Of Higher Rates Of Angle Class Iii Malocclusion Among Asian Populations . [42] Literature Also States That Previously Reported Angle Class Iii Prevalence Data Appears To Be Haphazard At Best Among Different Populations. Furthermore, Major Discrepancies Appear To Exist Among Populations Within Individual Countries. In The Limits Of Our Study Among Patients Who Reported To The Orthodontic Department For Treatment, 36.2% Of The Study Population Were Diagnosed With Class Iii Malocclusions. Only Patients With True Class Iii Malocclusion Were Included In The Study. 78% Of The Patients Were Diagnosed With A Skeletal Class Iii And 22% Of The Patients Were Diagnosed With A Class Iii Subdivision.(Table 1). The Results Of These And Our Study Are Not Comparable Because Our Study Accomplished Only On The Orthodontic Patient Population.

There Are Many Reasons Why Canines Fail To Erupt.[43] Most Surgeons Agree The Reasons May Include A Suspected Pathological Condition, Infection, Interference With Prosthetic Devices, Disturbance Of The Existing Dentition, Pain, And Ectopic Eruption. Many Authors Have Also Speculated About The Cause Of Impacted Canines And Their Relation To Malocclusion[6] . These Causes Include Inadequate Space, Supernumerary Teeth, Premature Loss Of The Deciduous Dentition, Retention Of The Deciduous Canine, Excessive Crown Length, Hereditary Factors, Functional Disturbances Of The Endocrine Glands, Tumors, Cysts, And Trauma. [44]Mitchell[45] Reported Trauma Also Has An Effect On The Impaction Of A Tooth As An Etiologic Factor.

In This Study,(Graph 1) The Incidence Of Canine Impaction In Men Was More Than Women (6% In Men, 2 % In Women), Being Statistically Significant($P<0.05$). Nagahara Et Al Observed That The Frequency Of Tooth Impaction Is Not Different Between Men And Women[46] . In Contrast, Dachi And Howell Et Al Stated That Girls Most Likely Have Canine Impaction Twice More Than Boys[47] . The Prevalence Of Impacted Maxillary Canines Is 0.9–2.2% In A Population With Normal Occlusion.[4] In The Limits Of Our Study , The Incidence Of Impacted Canine Among Patients With Class Iii Malocclusions Was Found Was 8% Of The Study Population Has Impacted Canines. Among The Study Population , Canine Impaction Showed More Association With Dental Class Iii Patients . A Chi-Square Analysis Done Among Patients With Class Iii Malocclusions And Of Canine Impaction , Gives A P Value Of 0.038 , So This Association Between Dental Class Iii Malocclusion And Canine Impaction Was Significant ($P> 0.05$). In Another Study, A Researcher Observed That The Prevalence Of Canine Impaction Was Not Different Significantly Between Patients With Class Iii Malocclusion [48] . Also, Leifert Et Al Reported That Angle C [49] . A Study Population With A Much Bigger And Unbiased Study Population May Help Shed A Light On The Association Between Malocclusions And Impacted Canine . Our Institution Is Passionate About High Quality Evidence Based Research And Has Excelled In Various Fields ([50–60].

Conclusion :

Within The Limited Sample Size, Canine Impaction Was Found In 8% Of Class Iii Malocclusion Patients. Canine Impaction Has Statistically Significant Association Among Both Adult And Child Age Groups, $P=0.028$. Also Males Patients With Class Iii Malocclusions Showed A Higher Association To Canine Impaction , $P=0.004$. Among The Study Population , Canine Impaction Showed More Association With Dental Class Iii Patients , $P = 0.038$.

Acknowledgment

The Authors Take Pleasure To Express Our Sincere Gratitude To The University For Granting Us Permission To Utilize The Data From Patient Records For The Study .

Authors Contribution

Author 1 (Amina Mehrin Bano) Carried Out The Retrospective Study By Collecting Data And Drafting The Manuscript After Performing The Necessary Statistical Analysis . Author 2(Dr.Srirengalakshimi) Aided In Conception Of The Topic , Has Participated In The Study Design , Statistical Analysis And Has Supervised In Preparation Of The Manuscript . All Authors Discussed The Results And Contributed To The Final Manuscript.

Conflict Of Interest

The Researcher Claims No Conflict Of Interest

Reference :

- [1] Richardson G, Russell Ka. A Review Of Impacted Permanent Maxillary Cuspids-Diagnosis And Prevention. *Journal-Canadian Dental Association* 2000;66:497–502.
- [2] Yavuz Ms, Aras Mh, Büyükkurt Mc. Impacted Mandibular Canines. *J Contemp Dent Pract* 2007.
- [3] D'amico Rm, Bjerklind K, Kurol J, Falahat B. Long-Term Results Of Orthodontic Treatment Of Impacted Maxillary Canines. *Angle Orthod* 2003;73:231–8.
- [4] Rajic S, Muretic Z, Percac S. Impacted Canine In A Prehistoric Skull. *Angle Orthod* 1996;66:477–80.
- [5] Bishara Se, Kommer Dd, Mcneil Mh, Montagano Ln, Oesterle Lj, Youngquist Hw. Management Of Impacted Canines. *Am J Orthod* 1976;69:371–87.
- [6] Smith Rj, Bailit Hl. Problems And Methods In Research On The Genetics Of Dental Occlusion. *Angle Orthod* 1977.
- [7] Abu-Hussein M, Sarafianou A. Mathematical Analysis Of Dental Arch Of Children In Normal Occlusion: A Literature Review. *International Journal Of Medical* 2012.
- [8] Abu-Hussein M, Watted N, Azzaldeen A. Prevalence Of Malocclusion And Impacted Canine In Arab Israelian Population (Arab48). *Journal Of Public ...* 2015.
- [9] Gill Ds, Naini Fb. *Orthodontics: Principles And Practice*. John Wiley & Sons; 2012.
- [10] Angle Eh. *Classification Of Malocclusion*. Dental Cosmos. St Louis 1899:248–64.
- [11] Proffit Wr, Fields Hw Jr, Sarver Dm. *Contemporary Orthodontics*. Elsevier Health Sciences; 2006.
- [12] Sivamurthy G, Sundari S. Stress Distribution Patterns At Mini-Implant Site During Retraction And Intrusion— A Three-Dimensional Finite Element Study. *Prog Orthod* 2016;17:4.
- [13] Krishnan S, Saravana Pandian Aks. Effect Of Bisphosphonates On Orthodontic Tooth Movement—An Update. *Journal Of Clinical And* 2015.
- [14] Vikram Nr, Prabhakar R, Kumar Sa, Karthikeyan Mk, Saravanan R. Ball Headed Mini Implant. *J Clin Diagn Res* 2017;11:Z102–3.
- [15] Viswanath A, Ramamurthy J, Dinesh Sps, Srinivas A. Obstructive Sleep Apnea: Awakening The Hidden Truth. *Niger J Clin Pract* 2015;18:1–7.
- [16] Felicita As. Quantification Of Intrusive/Retraction Force And Moment Generated During En-Masse Retraction Of Maxillary Anterior Teeth Using Mini-Implants: A Conceptual Approach. *Dental Press J Orthod* 2017;22:47–55.
- [17] Rubika J, Felicita As, Sivambiga V. Gonial Angle As An Indicator For The Prediction Of Growth Pattern. *World J Dent* 2015;6:161–3.
- [18] Jain Rk, Kumar Sp, Manjula Ws. Comparison Of Intrusion Effects On Maxillary Incisors Among Mini Implant

- Anchorage, J-Hook Headgear And Utility Arch. *J Clin Diagn Res* 2014.
- [19] Saravana Pandian K, Krishnan S, Aravind Kumar S. Angular Photogrammetric Analysis Of The Soft-Tissue Facial Profile Of Indian Adults. *Indian J Dent Res* 2018;29:137.
- [20] Ramesh Kumar Kr, Shanta Sundari Kk, Venkatesan A, Chandrasekar S. Depth Of Resin Penetration Into Enamel With 3 Types Of Enamel Conditioning Methods: A Confocal Microscopic Study. *Am J Orthod Dentofacial Orthop* 2011;140:479–85.
- [21] Dinesh Sps, Arun Av, Sundari Kks. An Indigenously Designed Apparatus For Measuring Orthodontic Force. *Journal Of Clinical And* 2013.
- [22] Felicita As, Chandrasekar S, Shanthasundari Kk. Determination Of Craniofacial Relation Among The Subethnic Indian Population: A Modified Approach-(Sagittal Relation). *Indian J Dent Res* 2012;23:305.
- [23] Samantha C, Sundari S, Chandrasekhar S. Comparative Evaluation Of Two Bis-Gma Based Orthodontic Bonding Adhesives-A Randomized Clinical Trial. *Journal Of Clinical And* 2017.
- [24] Kamisetty Sk, Verma Jk, Arun Ss. Sbs Vs Inhouse Recycling Methods-An Invitro Evaluation. *Journal Of Clinical And* 2015.
- [25] Felicita As. Orthodontic Management Of A Dilacerated Central Incisor And Partially Impacted Canine With Unilateral Extraction--A Case Report. *The Saudi Dental Journal* 2017;29:185–93.
- [26] Felicita As. Orthodontic Extrusion Of Ellis Class Viii Fracture Of Maxillary Lateral Incisor--The Sling Shot Method. *The Saudi Dental Journal* 2018;30:265–9.
- [27] Hafeez N, Others. Accessory Foramen In The Middle Cranial Fossa. *Research Journal Of Pharmacy And Technology* 2016;9:1880.
- [28] Krishnan Rp, Ramani P, Sherlin Hj, Sukumaran G, Ramasubramanian A, Jayaraj G, Et Al. Surgical Specimen Handover From Operation Theater To Laboratory: A Survey. *Ann Maxillofac Surg* 2018;8:234–8.
- [29] Somasundaram S, Ravi K, Rajapandian K, Gurunathan D. Fluoride Content Of Bottled Drinking Water In Chennai, Tamilnadu. *J Clin Diagn Res* 2015;9:Zc32.
- [30] Felicita As, Sumathi Felicita A. Orthodontic Extrusion Of Ellis Class Viii Fracture Of Maxillary Lateral Incisor – The Sling Shot Method. *The Saudi Dental Journal* 2018;30:265–9. <https://doi.org/10.1016/j.sdentj.2018.05.001>.
- [31] Kumar S, Rahman R. Knowledge, Awareness, And Practices Regarding Biomedical Waste Management Among Undergraduate Dental Students. *Asian J Pharm Clin Res* 2017;10:341.
- [32] Gurunathan D, Shanmugaavel Ak. Dental Neglect Among Children In Chennai. *J Indian Soc Pedod Prev Dent* 2016;34:364.
- [33] Sneha S, Others. Knowledge And Awareness Regarding Antibiotic Prophylaxis For Infective Endocarditis Among Undergraduate Dental Students. *Asian Journal Of Pharmaceutical And Clinical Research* 2016:154–9.
- [34] Dhinesh B, Isaac Joshuaramesh Lalvani J, Parthasarathy M, Annamalai K. An Assessment On Performance, Emission And Combustion Characteristics Of Single Cylinder Diesel Engine Powered By Cymbopogon Flexuosus Biofuel. *Energy Convers Manage* 2016;117:466–74.
- [35] Choudhari S, Thenmozhi Ms. Occurrence And Importance Of Posterior Condylar Foramen. *Laterality* 2016;8:11–43.
- [36] Paramasivam A, Vijayashree Priyadharsini J, Raghunandhakumar S. N6-Adenosine Methylation (M6a): A Promising New Molecular Target In Hypertension And Cardiovascular Diseases. *Hypertens Res* 2020;43:153–4.
- [37] Wu F, Zhu J, Li G, Wang J, Veeraraghavan Vp, Krishna Mohan S, Et Al. Biologically Synthesized Green Gold Nanoparticles From Siberian Ginseng Induce Growth-Inhibitory Effect On Melanoma Cells (B16). *Artif Cells Nanomed Biotechnol* 2019;47:3297–305.
- [38] Palati S, Ramani P, Shrelin H, Sukumaran G, Ramasubramanian A, Don Kr, Et Al. Knowledge, Attitude And Practice Survey On The Perspective Of Oral Lesions And Dental Health In Geriatric Patients Residing In Old Age Homes. *Indian Journal Of Dental Research* 2020;31:22. https://doi.org/10.4103/Ijdr.Ijdr_195_18.
- [39] Saravanan M, Arokiyaraj S, Lakshmi T, Pugazhendhi A. Synthesis Of Silver Nanoparticles From Phenerochaete Chrysosporium (Mtcc-787) And Their Antibacterial Activity Against Human Pathogenic Bacteria. *Microb Pathog* 2018;117:68–72.
- [40] Govindaraju L, Gurunathan D. Effectiveness Of Chewable Tooth Brush In Children-A Prospective Clinical Study. *J Clin Diagn Res* 2017;11:Zc31.
- [41] Vijayakumar Jain S, Muthusekhar Mr, Baig Mf, Senthilnathan P, Loganathan S, Abdul Wahab Pu, Et Al. Evaluation Of Three-Dimensional Changes In Pharyngeal Airway Following Isolated Lefort One Osteotomy For The Correction Of Vertical Maxillary Excess: A Prospective Study. *J Maxillofac Oral Surg* 2019;18:139–46.

Prevalence And Analysis Of Factors Associated With Canine Impaction In Class Iii Malocclusion. - A Retrospective Case Study


- [42] Hardy Dk, Cubas Yp, Orellana Mf. Prevalence Of Angle Class Iii Malocclusion: A Systematic Review And Meta-Analysis 2012.
- [43] Wright Dm. Forced Eruption Of An Impacted Lower Canine In A 48-Year-Old Man. *The Journal Of The American Dental Association* 1995.
- [44] Camilleri S, Scerri E. Transmigration Of Mandibular Canines—A Review Of The Literature And A Report Of Five Cases. *Angle Orthod* 2003.
- [45] Mitchell L. Displacement Of A Mandibular Canine Following Fracture Of The Mandible. *Br Dent J* 1993;174:417–8.
- [46] Nagahara K, Yuasa S, Yamada A, Ito K, Watanabe O, Iizuka T, Et Al. Etiological Study Of Relationship Between Impacted Permanent Teeth And Malocclusion. *Aichi Gakuin Daigaku Shigakkai Shi* 1989;27:913–24.
- [47] Dachi Sf, Howell Fv. A Survey Of 3,874 Routine Full-Mouth Radiographs: Ii. A Study Of Impacted Teeth. *Oral Surg Oral Med Oral Pathol* 1961.
- [48] Basdra Ek, Kiokpasoglou M, Stellzig A. The Class Ii Division 2 Craniofacial Type Is Associated With Numerous Congenital Tooth Anomalies. *Eur J Orthod* 2000;22:529–35.
- [49] Leifert S, Jonas Ie. Dental Anomalies As A Microsymptom Of Palatal Canine Displacement. *J Orofac Orthop* 2003;64:108–20.
- [50] Vijayashree Priyadharsini J. In Silico Validation Of The Non-Antibiotic Drugs Acetaminophen And Ibuprofen As Antibacterial Agents Against Red Complex Pathogens. *J Periodontol* 2019;90:1441–8.
- [51] Pc J, Marimuthu T, Devadoss P. Prevalence And Measurement Of Anterior Loop Of The Mandibular Canal Using Cbct: A Cross Sectional Study. *Clin Implant Dent Relat Res* 2018.
- [52] Ramesh A, Varghese S, Jayakumar Nd, Malaiappan S. Comparative Estimation Of Sulfiredoxin Levels Between Chronic Periodontitis And Healthy Patients - A Case-Control Study. *J Periodontol* 2018;89:1241–8.
- [53] Ramadurai N, Gurunathan D, Samuel Av, Subramanian E, Rodrigues Sjl. Effectiveness Of 2% Articaine As An Anesthetic Agent In Children: Randomized Controlled Trial. *Clin Oral Investig* 2019;23:3543–50.
- [54] Sridharan G, Ramani P, Patankar S, Vijayaraghavan R. Evaluation Of Salivary Metabolomics In Oral Leukoplakia And Oral Squamous Cell Carcinoma. *J Oral Pathol Med* 2019;48:299–306.
- [55] Ezhilarasan D, Apoorva Vs, Ashok Vardhan N. Syzygium Cumini Extract Induced Reactive Oxygen Species-Mediated Apoptosis In Human Oral Squamous Carcinoma Cells. *J Oral Pathol Med* 2019;48:115–21.
- [56] Mathew Mg, Samuel Sr, Soni Aj, Roopa Kb. Evaluation Of Adhesion Of Streptococcus Mutans, Plaque Accumulation On Zirconia And Stainless Steel Crowns, And Surrounding Gingival Inflammation In Primary Molars: Randomized Controlled Trial. *Clin Oral Investig* 2020:1–6.
- [57] Samuel Sr. Can 5-Year-Olds Sensibly Self-Report The Impact Of Developmental Enamel Defects On Their Quality Of Life? *Int J Paediatr Dent* 2021;31:285–6.
- [58] R H, Hannah R, Ramani P, Ramanathan A, R Jm, Gheena S, Et Al. Cyp2 C9 Polymorphism Among Patients With Oral Squamous Cell Carcinoma And Its Role In Altering The Metabolism Of Benzo[A]Pyrene. *Oral Surgery, Oral Medicine, Oral Pathology And Oral Radiology* 2020;130:306–12. <https://doi.org/10.1016/j.oooo.2020.06.021>.
- [59] Chandrasekar R, Chandrasekhar S, Sundari Kks, Ravi P. Development And Validation Of A Formula For Objective Assessment Of Cervical Vertebral Bone Age. *Prog Orthod* 2020;21:38.
- [60] Vijayashree Priyadharsini J, Smiline Girija As, Paramasivam A. In Silico Analysis Of Virulence Genes In An Emerging Dental Pathogen *A. Baumannii* And Related Species. *Arch Oral Biol* 2018;94:93–8.

Tables And Graphs :

	Presence Of Impacted Canine	
	Impacted Canine	No Impacted Canine


			Number Of Patients (%)	Number Of Patients (%)
Type Of Malocclusion	Skeletal Class Iii	Male	2(4%)	20(40%)
		Female	0(0%)	17(34%)
	Dental Class Iii	Male	1(2%)	4(8%)
		Female	1(2%)	5(10%)

Table 1: Table Describes The Study Population Based On Gender Anf Type Of Class Iii Malocclusion . Canine Impaction Is Seen In 4% Of The Male Population With Skeletal Class Iii Malocclusion And 2% Of The Dental Class Iii Malocclusions .Also , Among The Female Study Population Canine Malocclusion Is Seen In Only 2% Of The Dental Class Iii Malocclusions And None In Skeletal Class Iii Malocclusion .




Prevalence And Analysis Of Factors Associated With Canine Impaction In Class Iii Malocclusion. - A Retrospective Case Study

Graph 1 : Graph Depicts The Association Between Presence Of Impacted Canine Patients Diagnosed With Class Iii Malocclusion And Gender. The X Axis Denotes The Gender Of The Patients And The Y Axis Denotes The Number Of Patients With Or Without Impacted Canine. Among The Study Population , Male Patients Were Found To Have 6% Of Impacted Canine(Blue) . The Female Study Patients Were Found To Have 2% Of Impacted Canine(Blue). Chi Square Test. $P=0.004$,(<0.05) Hence Statistically Significant .



Graph 2: Graph Depicts An Association Between The Type Of Malocclusion And Presence Of Impacted Canine .The X Axis Represents The Type Of Class Iii Malocclusion And The Y Axis Represents The Number Of Patients With Or Without Canine Impactions .Among The Patients With Skeletal Class Iii , 2% Of The Patients Have Impacted Canines (Blue). Also Among Patients With Dental Class Iii, 2% Of The Patients Have Impacted Canines (Blue) .Impacted Canines Have More Association With Dental Class Iii Patients ,Chi-Square Test Done , Chi Square Test Done. $P = 0.038$ (<0.05) Which Is Statistically Significant.



Graph 3: Graph Shows The Association Between Incidence Of Impacted Canine And Age. The X-Axis Represents Incidence Of Impacted Canine And Y-Axis Represents The Number Of Patients . About 4% Of The Study Population With Impacted Canines Belong To The Adult Age Group (Green) And 4% Of The Study Population With Impacted Canines Belong To The Age Agroung Of Children (Blue)Chi Square Test Done. $P=0.028$ (<0.05) , Hence Statistically Significant .