

## **Recurrence of Oral cancer following treatment in an institutional set up - A retrospective analysis of the associated factor.**

**Sindhu Priya Kuppusamy Sundara Murthy**

Saveetha Dental College and Hospitals,  
Saveetha Institute of Medical and Technical sciences (SIMATS),  
Saveetha University  
Chennai - 77, India.  
Email ID: 151501024.sdc@saveetha.com  
Contact number: +91 7708045340

**Dr. Mahathi. N**

Reader,  
Department of Oral and Maxillofacial surgery,  
Saveetha Dental College and Hospitals,  
Saveetha Institute of Medical and Technical sciences (SIMATS),  
Saveetha University  
Chennai - 77, India.  
Email ID: mahathin.sdc@saveetha.com

**Dr. Suresh**

Reader,  
Department of Prosthodontics,  
Saveetha Dental College and Hospitals,  
Saveetha Institute of Medical and Technical sciences (SIMATS),  
Saveetha University  
Chennai - 77, India.  
Email ID: suresh@saveetha.com

### **Abstract**

Incidence of cancers of the oral cavity is increasing and has replaced laryngeal cancer as the most frequently occurring cancer in the head and neck area. Routine follow up is must for a period of 6 months to 1 year after treatment. Current guidelines advocate a follow-up period of at least 5 years to evaluate the time of recurrence following the treatment of oral cancer. Data samples required for study were taken from hospital records. All the collected data were cross verified and compiled together in an excel sheet. Compiled data were statistically analysed in SPSS software. In this study, a total of 51 patients with oral cancer had undergone treatment. 74.51% of patients reported back without any recurrence after a follow-up period of 6 months to 1 year. On associating recurrence with age it was noted that the recurrence was more in higher age groups p value of 0.025. Recurrence was common in males than females with p value of 0.025 (<0.05). To conclude, the majority of the patients in the study had no recurrence following treatment in the above follow-up period.

**Keywords:** Laryngeal cancer, Recurrence, Tumour.

### **Introduction**

Head and neck squamous cell carcinoma are the 6<sup>th</sup> most common malignancies in the world. The oral cavity is the most common subsite and makes upto 3% of all cancer cases worldwide. Oral squamous cell carcinoma has poor prognosis and in case of relapse with mortality rates upto 92%<sup>1,2,3</sup>.

Surgery is the preferred treatment for squamous cell carcinoma. Despite great progress in chemotherapy and radiotherapy, the prognosis of OSCC is poor due to aggressive local invasion and metastasis, leading to recurrence<sup>4</sup>.

After curative treatment of primary tumour, recurrence rates range from 7 - 47.7% have been reported. Even though significant improvements in diagnostics and treatment modalities were made, tumour relapse is one of the major problems. Upto 86% of all recurrent tumours recur in the first 2 years after treatment<sup>5,6</sup>.

## Recurrence of Oral cancer following treatment in an institutional set up - A retrospective analysis of the associated factor.

The incidence, recurrence and mortality rate of OCC are increasing. Monitoring the trends in OCC is essential for optimised prevention, treatment, and surveillance care for the growing group of patients, therefore in order to slightly improve the survival rates<sup>7</sup>. Previously our team has a rich experience in working on various research projects across multiple disciplines The (1–3)(4–15)

The aim of this study is to evaluate the time of recurrence following the treatment of oral cancer in a private hospital.

### Materials and method

All the data of patients who underwent treatment for oral cancer and returned with recurrence of oral cancer were taken for the study as a sample. The study setting was a university setting. Exclusion criteria was case sheets with incomplete data and those patients who did not come for follow up visits when called.

Data was collected from case sheets of patients who reported during the months of October 2019 and March 2020 from the hospital record management system where all the records of patients regarding their medical and dental history and treatment done are stored. Cross verification was done to avoid bias by another examiner. To avoid missing any data, photographic evaluation was done.

All the relevant data were retrieved and tabulated in excel sheet. Later, it was statistically analysed by IBM SPSS statistics, using the Chi-Square test. Independent variables are the total number of treated patients and dependent variables are recurrence of oral cancer.

### Results

The study consisted of a total of 51 patients, among which 34 patients were males, and 17 patients were females. All the 51 patients had undergone treatment for oral cancer, among which 12 patients had recurrence of oral cancer and 39 had no recurrence. The percentage of patients with recurrence is 25.49% and without recurrence is 74.51% (Fig.1). 5.88% from 46–60 and 19.61% from >60 age groups had recurrence. 19.61% from 30–45, 21.57% from 46–60 and 33.33% from >60 age groups had no recurrence with p value of 0.025 (<0.05) (Fig.3). 15.69% of males had recurrence and the rest 56.86% had no recurrence. The number of females who had recurrence of oral cancer were 9.80% and without was 17.65% with p value of 0.025 (<0.05) (Fig.2). The percentage of recurrence that occurred within 1–2 months was 58.33%, within 3–4 months was 25% and after more than 5 months was 15.62% (Fig.4).

### Discussion

In the present study, a higher percentage of patients did not have recurrence of oral cancer.

In a study by Bo wang, 32.7% of patients had recurrence. Chi square test showed that in the T-stage, degrees of differentiation were important factors of recurrence. The 2- and 5- year survival rates were lower in patients with recurrence than in those without<sup>23,24</sup>. Factors that influence the recurrence of OSCC have been explored in recent years. Ebrahim et al and Camisasca have analysed patient's clinico pathological data, including tumour sites, clinical and pathological stages, invasion mode and perineural invasion. They concluded that tongue cancer and poor differentiation contributed to recurrence after surgery<sup>25</sup>.

Vasquez et Al reported recurrence rate of 44.9% in 118 patients. The important factors were co-morbidities rate, degree of tumour differentiation and stage. In another study, it suggested that when the treatment involved only surgery, the recurrence rate was higher. When combined with chemotherapy and radiotherapy, complete tumour resection with various flaps reduces tumour recurrence, suggested by Bo Wang<sup>26,27</sup>.

Vincente et Al followed up to 98 patients with OSCC, the mortality rate 47% in patients with flap repair and 67% in patients without flap repair. Capote et Al suggested careful removal of lymph nodes with primary resection of tumours. Because the OSCC might metastasize into the sternocleidomastoid muscle lymph groups, these should be removed during surgery. The recurrence rate in their study was 32.7%. Lopez Rodriguez et al reported that preoperative radiotherapy and chemotherapy for OSCC at N2-N3 stage can completely control neck lymph node metastasis<sup>28,29</sup>.

In the present study, the reason for the low recurrence rate of 25.49% is highly due to combination therapy of surgery and chemotherapy and radiotherapy. Our institution is passionate about high quality evidence based research and has excelled in various fields ( (16–26)

This study could be further improved by increasing the sample size and checking at a molecular level using biomarkers to predict the recurrence and prognosis of oral cancer.

### Conclusion

Oral squamous cell carcinoma is a commonly occurring oral cancer and it is associated with significant mortality and morbidity. The purpose of this study was to evaluate the recurrence of oral cancer following treatment in order to provide better prognosis and quality of life. In our study, the majority of the patients did not have recurrence and it was found to be statistically significant that the majority of people had recurrence within 1-2 months following treatment.

**AUTHORS CONTRIBUTION** First author, Sindhupriya performed the data collection by reviewing patient details, filtering required data, analysing and interpreting statistics and contributed to manuscript writing.

Second author, Dr. Mahathi contributed to conception of study title, study design, analysed the collected data, statistics and interpretation and also critically revised the manuscript.

Third author, Dr. Suresh. V participated in the study and revised the manuscript. All the three authors have discussed the results and contributed to the final manuscript.

**ACKNOWLEDGMENT** This research was supported by Saveetha Dental College and Hospital. We thank the department of Oral Medicine, Saveetha Dental College for providing insight and expertise that greatly assisted this research.

**CONFLICT OF INTEREST** The authors declare that there is no conflict of interests.

### References

1. Bagan JV, Scully C. Recent advances in Oral Oncology 2007: epidemiology, etiopathogenesis, diagnosis and prognostication. *Oral Oncol* [Internet]. 2008 Feb;44(2):103–8. Available from: <http://dx.doi.org/10.1016/j.oraloncology.2008.01.008>
2. van Dijk BAC, Brands MT, Geurts SME, Merckx MAW, Roodenburg JLN. Trends in oral cavity cancer incidence, mortality, survival and treatment in the Netherlands. *Int J Cancer* [Internet]. 2016 Aug 1;139(3):574–83. Available from: <http://dx.doi.org/10.1002/ijc.30107>
3. Rogers SN, Brown JS, Woolgar JA, Lowe D, Magennis P, Shaw RJ, et al. Survival following primary surgery for oral cancer. *Oral Oncol* [Internet]. 2009 Mar;45(3):201–11. Available from: <http://dx.doi.org/10.1016/j.oraloncology.2008.05.008>
4. Johnson FE, Maehara Y, Browman GP, Margenthaler JA, Audisio RA, Thompson JF, et al. Patient Surveillance After Cancer Treatment [Internet]. Springer Science & Business Media; 2013. 538 p. Available from: <https://play.google.com/store/books/details?id=ZcZDAAAQBAJ>
5. McGarry GW, Mackenzie K, Periasamy P, McGurk F, Gatehouse S. Multiple primary malignant tumours in patients with head and neck cancer: the implications for follow-up. *Clin Otolaryngol Allied Sci* [Internet]. 1992;17(6):558–62. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1365-2273.1992.tb01718.x>
6. Curtis RE. New Malignancies Among Cancer Survivors: SEER Cancer Registries, 1973-2000 [Internet]. U.S. Department of Health and Human Services, National Institutes of Health, National Cancer Institute; 2006. 492 p. Available from: <https://play.google.com/store/books/details?id=z28dJN3EdAYC>
7. Coviello V, Boggess M. Cumulative Incidence Estimation in the Presence of Competing Risks. *Stata J* [Internet]. 2004 Jun 1;4(2):103–12. Available from: <https://doi.org/10.1177/1536867X0400400201>
8. Jesudasan JS, Wahab PA, Sekhar MM. Effectiveness of 0.2% chlorhexidine gel and a eugenol-based paste on postoperative alveolar osteitis in patients having third molars extracted: a randomised controlled clinical trial. *British Journal of Oral and Maxillofacial Surgery*. 2015 Nov 1;53(9):826-30.
9. Santhosh Kumar, M.P., RAHMAN R. Knowledge, awareness, and practices regarding biomedical waste management among undergraduate dental students. *Asian J Pharm Clin Res*. 2017;10(8):341- 5.
10. Christabel A, Anantanarayanan P, Subash P, Soh CL, Ramanathan M, Muthusekhar MR, Narayanan V. Comparison of pterygomaxillary dysjunction with tuberosity separation in isolated Le Fort I osteotomies:

Recurrence of Oral cancer following treatment in an institutional set up - A retrospective analysis of the associated factor.

- A prospective, multi-centre, triple-blind, randomized controlled trial. International journal of oral and maxillofacial surgery. 2016 Feb 1;45(2): 180-5.
11. Marimuthu M, Andiappan M, Wahab A, Muthusekhar MR, Balakrishnan A, Shanmugam S. Canonical Wnt pathway gene expression and their clinical correlation in oral squamous cell carcinoma. Indian Journal of Dental Research. 2018 May 1;29(3):291.
  12. Packiri S, Gurunathan D, Selvarasu K. Management of paediatric oral ranula: a systematic review. Journal of clinical and diagnostic research: JCDR. 2017 Sep;11 (9):ZE06
  13. Santhosh Kumar, M.P. Relationship between dental anxiety and pain experience during dental extractions. Asian J Pharm Clin Res. 2017;10(3):458- 61.
  14. Patil SB, Durairaj D, Kumar GS, Karthikeyan D, Pradeep D. Comparison of extended nasolabial flap versus buccal fat pad graft in the surgical management of oral submucous fibrosis: a prospective pilot study. Journal of maxillofacial and oral surgery. 2017 Sep 1;16(3):312-21.
  15. Rao TD, Santhosh Kumar, M.P. Analgesic Efficacy of Paracetamol Vs Ketorolac after Dental Extractions. Research Journal of Pharmacy and Technology. 2018;11(8): 3375-9.
  16. Abhinav RP, Selvarasu K, Maheswari GU, Taltia AA. The patterns and etiology of maxillofacial trauma in South India. Annals of maxillofacial surgery. 2019 Jan;9(1): 114.
  17. Santhosh Kumar, M.P., Snea S. Knowledge and awareness regarding antibiotic prophylaxis for infective endocarditis among undergraduate dental students. Asian J Pharm Clin Res. 2016;9: 154-9.
  18. Santhosh Kumar, M.P. The emerging role of botulinum toxin in the treatment of orofacial disorders: Literature update. Asian J Pharm Clin Res. 2017;10(9):21-9.
  19. Rahman R, Santhosh Kumar, M.P. Knowledge, attitude, and awareness of dental undergraduate students regarding human immunodeficiency virus/acquired immunodeficiency syndrome patients. Asian J Pharm Clin Res. 2017;10(5):175- 80.
  20. Santhosh Kumar, M.P., Sneha S. Knowledge and awareness regarding antibiotic prophylaxis for infective endocarditis among undergraduate dental students. Asian J Pharm Clin Res. 2016;9: 154-9.
  21. Patturaja K, Pradeep D. Awareness of Basic Dental Procedure among General Population. Research Journal of Pharmacy and Technology. 2016;9(9): 1349-51.
  22. Jain SV, Muthusekhar MR, Baig MF, Senthilnathan P, Loganathan S, Wahab PA, Madhulakshmi M, Vohra Y. Evaluation of three-dimensional changes in pharyngeal airway following isolated lefort one osteotomy for the correction of vertical maxillary excess: a prospective study. Journal of maxillofacial and oral surgery. 2019 Mar 8;18(1):139-46.




Fig.1 Bar graph represents the percentage of patients with and without recurrence of cancer. X-axis represents the patients with and without recurrence and Y-axis represents the number of the patients. From the graph is evident that the percentage of patients with recurrence was 25.49% (yellow bar) and without recurrence was 74.51% (brown bar).




Fig.2 Bar graphs represent the association between gender and recurrence of oral cancer after treatment. X-axis represents the gender of patients. Y-axis represents the number of patients either recurrence after treatment. More males (15.69%) than females (9.80%) had recurrence of oral cancer (yellow bar). Chi-square test was done and the association was found to be statistically significant. Pearson's Chi-square value: 0.303, DF:2, p value: 0.025(<0.05), proving that more males had recurrence than females.




Fig.3 Bar graphs represent the association between age and recurrence of oral cancer after treatment. X-axis represents the age of patients. Y-axis represents the number of patients with recurrence. Majority of patients who had recurrence of oral cancer (19.61%) were above 60 years of age. Chi-square test was done and the association was found to be statistically significant. Pearson's Chi-square value: 0.327, DF:2, p value:

Recurrence of Oral cancer following treatment in an institutional set up - A retrospective analysis of the associated factor.

0.025(<0.05), proving that recurrence of oral cancer is common in older age groups.




Fig.4 Bar graph represents the time period of recurrence and the number of patients in each category. X-axis represents the time period of recurrence in months. Y-axis represents the number of the patients with recurrence. The percentage of patients with recurrence within 1-2 months is 58.33% (pink bar), within 3-4 months is 25% (green bar) and more than 5 months is 16.67% (purple bar).

Legend of graphs

Graph 1	Percentage of patients with recurrence of cancer
Graph 2	Gender association with recurrence of cancer
Graph 3	Age association with recurrence of cancer
Graph 4	Time of recurrence in months

1. Hafeez N, Others. Accessory foramen in the middle cranial fossa. *Research Journal of Pharmacy and Technology* [Internet]. 2016;9(11):1880. Available from: <https://search.proquest.com/openview/3790b1bd527fc74251e9798d8ab23bc8/1?pq-origsite=scholar&cbl=1096441>
2. Krishnan RP, Ramani P, Sherlin HJ, Sukumaran G, Ramasubramanian A, Jayaraj G, et al. Surgical Specimen Handover from Operation Theater to Laboratory: A Survey. *Ann Maxillofac Surg* [Internet]. 2018 Jul;8(2):234–8. Available from: [http://dx.doi.org/10.4103/ams.ams\\_51\\_18](http://dx.doi.org/10.4103/ams.ams_51_18)
3. Somasundaram S, Ravi K, Rajapandian K, Gurunathan D. Fluoride content of bottled drinking water in Chennai, Tamilnadu. *J Clin Diagn Res* [Internet]. 2015;9(10):ZC32. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4625331/>
4. Felicita AS, Sumathi Felicita A. Orthodontic extrusion of Ellis Class VIII fracture of maxillary lateral incisor – The sling shot method [Internet]. Vol. 30, *The Saudi Dental Journal*. 2018. p. 265–9. Available from: <http://dx.doi.org/10.1016/j.sdentj.2018.05.001>

5. Kumar S, Rahman R. Knowledge, awareness, and practices regarding biomedical waste management among undergraduate dental students. *Asian J Pharm Clin Res* [Internet]. 2017 Aug 1;10(8):341. Available from: <https://innovareacademics.in/journals/index.php/ajpcr/article/view/19101>
6. Gurunathan D, Shanmugaavel AK. Dental neglect among children in Chennai. *J Indian Soc Pedod Prev Dent* [Internet]. 2016 Oct 1 [cited 2020 Jun 8];34(4):364. Available from: <http://www.jisppd.com/article.asp?issn=0970-4388;year=2016;volume=34;issue=4;spage=364;epage=369;aulast=Gurunathan>
7. Sneha S, Others. Knowledge and awareness regarding antibiotic prophylaxis for infective endocarditis among undergraduate dental students. *Asian Journal of Pharmaceutical and Clinical Research* [Internet]. 2016;154–9. Available from: <https://innovareacademics.info/journals/index.php/ajpcr/article/download/13405/8142>
8. Dhinesh B, Isaac JoshuaRamesh Lalvani J, Parthasarathy M, Annamalai K. An assessment on performance, emission and combustion characteristics of single cylinder diesel engine powered by Cymbopogon flexuosus biofuel. *Energy Convers Manage* [Internet]. 2016 Jun 1;117:466–74. Available from: <https://www.sciencedirect.com/science/article/pii/S0196890416301868>
9. Choudhari S, Thenmozhi MS. Occurrence and Importance of Posterior Condylar Foramen. *Laterality* [Internet]. 2016;8:11–43. Available from: [https://rjptonline.org/HTML\\_Papers/Research%20Journal%20of%20Pharmacy%20and%20Technology\\_\\_PID\\_\\_2016-9-8-15.html](https://rjptonline.org/HTML_Papers/Research%20Journal%20of%20Pharmacy%20and%20Technology__PID__2016-9-8-15.html)
10. Paramasivam A, Vijayashree Priyadharsini J, Raghunandhakumar S. N6-adenosine methylation (m6A): a promising new molecular target in hypertension and cardiovascular diseases. *Hypertens Res* [Internet]. 2020 Feb;43(2):153–4. Available from: <http://dx.doi.org/10.1038/s41440-019-0338-z>
11. Wu F, Zhu J, Li G, Wang J, Veeraraghavan VP, Krishna Mohan S, et al. Biologically synthesized green gold nanoparticles from Siberian ginseng induce growth-inhibitory effect on melanoma cells (B16). *Artif Cells Nanomed Biotechnol* [Internet]. 2019 Dec;47(1):3297–305. Available from: <http://dx.doi.org/10.1080/21691401.2019.1647224>
12. Palati S, Ramani P, Shrelin H, Sukumaran G, Ramasubramanian A, Don KR, et al. Knowledge, Attitude and practice survey on the perspective of oral lesions and dental health in geriatric patients residing in old age homes [Internet]. Vol. 31, *Indian Journal of Dental Research*. 2020. p. 22. Available from: [http://dx.doi.org/10.4103/ijdr.ijdr\\_195\\_18](http://dx.doi.org/10.4103/ijdr.ijdr_195_18)
13. Saravanan M, Arokiyaraj S, Lakshmi T, Pugazhendhi A. Synthesis of silver nanoparticles from *Phenerochaete chrysosporium* (MTCC-787) and their antibacterial activity against human pathogenic bacteria. *Microb Pathog* [Internet]. 2018 Apr;117:68–72. Available from: <http://dx.doi.org/10.1016/j.micpath.2018.02.008>
14. GovinDaraju L, Gurunathan D. Effectiveness of Chewable Tooth Brush in Children-A Prospective Clinical Study. *J Clin Diagn Res* [Internet]. 2017;11(3):ZC31. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5427431/>
15. Vijayakumar Jain S, Muthusekhar MR, Baig MF, Senthilnathan P, Loganathan S, Abdul Wahab PU, et al. Evaluation of Three-Dimensional Changes in Pharyngeal Airway Following Isolated Lefort One Osteotomy for the Correction of Vertical Maxillary Excess: A Prospective Study. *J Maxillofac Oral Surg* [Internet]. 2019 Mar;18(1):139–46. Available from: <http://dx.doi.org/10.1007/s12663-018-1113-4>
16. Vijayashree Priyadharsini J. In silico validation of the non-antibiotic drugs acetaminophen and ibuprofen as antibacterial agents against red complex pathogens. *J Periodontol* [Internet]. 2019 Dec;90(12):1441–8. Available from: <http://dx.doi.org/10.1002/JPER.18-0673>
17. Pc J, Marimuthu T, Devadoss P. Prevalence and measurement of anterior loop of the mandibular canal using CBCT: A cross sectional study. *Clin Implant Dent Relat Res* [Internet]. 2018; Available from: <https://europepmc.org/article/med/29624863>

Recurrence of Oral cancer following treatment in an institutional set up - A retrospective analysis of the associated factor.

18. Ramesh A, Varghese S, Jayakumar ND, Malaiappan S. Comparative estimation of sulfiredoxin levels between chronic periodontitis and healthy patients - A case-control study. *J Periodontol* [Internet]. 2018 Oct;89(10):1241–8. Available from: <http://doi.wiley.com/10.1002/JPER.17-0445>
19. Ramadurai N, Gurunathan D, Samuel AV, Subramanian E, Rodrigues SJL. Effectiveness of 2% Articaine as an anesthetic agent in children: randomized controlled trial. *Clin Oral Investig* [Internet]. 2019 Sep;23(9):3543–50. Available from: <http://dx.doi.org/10.1007/s00784-018-2775-5>
20. Sridharan G, Ramani P, Patankar S, Vijayaraghavan R. Evaluation of salivary metabolomics in oral leukoplakia and oral squamous cell carcinoma. *J Oral Pathol Med* [Internet]. 2019 Apr;48(4):299–306. Available from: <http://dx.doi.org/10.1111/jop.12835>
21. Ezhilarasan D, Apoorva VS, Ashok Vardhan N. Syzygium cumini extract induced reactive oxygen species-mediated apoptosis in human oral squamous carcinoma cells. *J Oral Pathol Med* [Internet]. 2019 Feb;48(2):115–21. Available from: <http://dx.doi.org/10.1111/jop.12806>
22. Mathew MG, Samuel SR, Soni AJ, Roopa KB. Evaluation of adhesion of *Streptococcus mutans*, plaque accumulation on zirconia and stainless steel crowns, and surrounding gingival inflammation in primary molars: Randomized controlled trial. *Clin Oral Investig* [Internet]. 2020;1–6. Available from: <https://link.springer.com/article/10.1007/s00784-020-03204-9>
23. Samuel SR. Can 5-year-olds sensibly self-report the impact of developmental enamel defects on their quality of life? *Int J Paediatr Dent* [Internet]. 2021 Mar;31(2):285–6. Available from: <http://dx.doi.org/10.1111/ipd.12662>
24. R H, Hannah R, Ramani P, Ramanathan A, R JM, Gheena S, et al. CYP2 C9 polymorphism among patients with oral squamous cell carcinoma and its role in altering the metabolism of benzo[a]pyrene [Internet]. Vol. 130, *Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology*. 2020. p. 306–12. Available from: <http://dx.doi.org/10.1016/j.oooo.2020.06.021>
25. Chandrasekar R, Chandrasekhar S, Sundari KKS, Ravi P. Development and validation of a formula for objective assessment of cervical vertebral bone age. *Prog Orthod* [Internet]. 2020 Oct 12;21(1):38. Available from: <http://dx.doi.org/10.1186/s40510-020-00338-0>
26. Vijayashree Priyadharsini J, Smiline Girija AS, Paramasivam A. In silico analysis of virulence genes in an emerging dental pathogen *A. baumannii* and related species. *Arch Oral Biol* [Internet]. 2018 Oct;94:93–8. Available from: <http://dx.doi.org/10.1016/j.archoralbio.2018.07.001>