

Turkish Online Journal of Qualitative Inquiry (TOJQI)

Volume 5, July 2021 : 2908- 2917

Research Article

A Clinical Study on Anti-wrinkle Efficacy of Cosmetics Containing Oligoarginine Conjugation of Palmitoyl-GHK

Su In Park¹, Kwang Won Lee², Hoon Cha³, Moon Sam Shin^{4*}

Abstract

In this study, Arg 4 (R4) was conjugated with Pal-GHK to get palmitoyl heptapeptide, Pal-GHK-R4 and wrinkle improvement clinical trials of cosmetics containing Pal-GHK-R4 were conducted. As a result of total 21 subjects used the test products 4 weeks use on the face area; 1) measurement results of crow's feet after 2 and 4 weeks of test products use, analysis value of crow's feet statistically significantly decreased after 2 and 4 weeks use compared to before use of products ($p<0.05$); 2) measurement results of skin moisturization after 2 and 4 weeks test products use, measured value of skin moisture content statistically significantly increased after 2 and 4 weeks use compared to before use of products ($p<0.05$); 3) measurement results of transepidermal water loss (TEWL) after 2 and 4 weeks test products use, measured value of TEWL statistically significantly decreased after 2 and 4 weeks use compared to before use of products ($p<0.05$); 4) there was no skin adverse event reported after using the products during the study period. Therefore, the test products containing both wrinkle improving and skin penetrating peptides, Pal-GHK-R4 are considered to have beneficial effects on improvement of skin wrinkle, moisturization, TEWL after 2 and 4 weeks use of test products.

Keywords: *Cell penetrating peptides, transdermal delivery, conjugated liposomes, skin permeation.*

¹ Ph.D. Student, Eulji University, Senior Healthcare majoring in Cosmetic Pharmacology, sooo_30@naver.com

² Master Student, Eulji University, Senior Healthcare majoring in Cosmetic Pharmacology, eecl308@naver.com

³ CEO, Dermafirm Co. Ltd, R&D Center, ceo@dermafirm.kr.

^{4*} Professor, Eulji University, Senior Healthcare majoring in Cosmetic Pharmacology, msshin@eulji.ac.kr

Introduction

The aging population structure is not limited to developed countries but is a phenomenon that is being carried out worldwide including developing countries. The increase of the elderly causes various social problems, but also leads to the growth of the senior industry (Yang, 2010). In the cosmetics market, senior generation women have emerged as a high consumer and a high purchasing power, and they are a big part of important consumers (Oh, 2017). As the number of senior generation women who want young and resilient skin increases, the demand for functional cosmetics that improve wrinkles with excellent efficacy has increased, and the cosmetics industry is actively pursuing research on this (Kang, 2007).

Recently, there has been a growing interest in using peptides, an amino acid polymer, as a functional material for wrinkle improvement in cosmetics. Peptides have excellent physiological activity, are safe because they are *in vivo*, are stable in light and air, and are smaller than proteins, so they can be synthesized, so they are highly likely to be used as cosmetic materials (Moh, 2011; Yoo, 2017). GHK (glycyl-histidyl-lysine), which was first isolated from human serum and known as a hepatocellular growth factor, is a typical wrinkle-improving functional peptide. GHK exists in the blood about 200 mg/mL at the age of 20, but it decreases as aging progresses and is about 80 mg/mL at the age of 60. This increases the synthesis of extracellular matrix (ECM) such as collagen, elastin, and glycosaminoglycans (GAGs), which can cause skin regeneration and wound healing. GHK is commonly present in the form of GHK-Cu with high affinity to copper ions, and Pal-GHK, which is bound with palmitate, a fatty acid, is also present to enhance the permeability and stability of GHK. These GHK, GHK-Cu, and Pal-GHK are popular as functional cosmetic materials for wrinkle improvement. However, the problem is that the peptide has difficulty in penetrating the stratum corneum due to its hydrophilic properties, making it difficult to actually exhibit efficacy in the skin (Li, 2015; Maquart, 1988; Pickart, 2008; Maquart, 1993; Godet, 1995; Pickart, 1980).

On the other hand, there have been active studies to deliver hydrophilic macromolecules without destroying the cell membrane and without showing cytotoxicity for effective intracellular drug delivery. In this regard, many studies on cell penetrating peptides (CPPs) have been conducted. Cell penetrating peptides are generally composed of 10-30 short amino acids, and most contain a large number of basic amino acids, lysine (K) and arginine (R). TAT peptides derived from HIV-1 are one of the most studied cell penetrating peptides. TAT peptides are composed of 86 amino acids, among which the sequence of RKKKRRQRRR is known to be the minimum site for cell penetrating peptides. The experiments that substituted

or deleted amino acid sequences of TAT peptides revealed that lysine (K) and arginine (R) play a key role in the intracellular permeability of TAT peptides as cell penetrating peptides (Lee, 2019; Vive, 1997; Wang, 2019; Yi, 2007; Liang, 2004). Furthermore, since both the cell membrane and intercellular lipids have structural similarity in the structure of lipid bilayer, it is considered that the cell penetrating peptides can be applied to the intercellular lipids' permeation of functional materials (Mazurowska, 2008; Kim, 2009).

In this study, Arg 4 (R4) was conjugated with Pal-GHK to get palmitoyl heptapeptide, Pal-GHK-R4 and clinical study on improvement of crow's feet, skin moisturization, transepidermal water loss (TEWL) cosmetics containing Pal-GHK-R4 were conducted.

Materials and Methods

Study Protocols and Measurements

The cosmetics used in this test are "Biotoc Regen" manufactured by Dermafirm Co., Ltd., which consists of 4 products (ampoule, serum, cream and peel). The main ingredients contained 1,000 ppm (0.1%) of Pal-GHK-R4. The other ingredients contained 99.9% of emulsifier, oil, humectants, fragrance and deionized water. Pal-GHK-R4 is manufactured from Dermafirm Co., Ltd., in Korea and have a purity of at least 99.0%, respectively.

In this study, KC Skin Research Center conducted the body efficacy evaluation according to the tenets of the Declaration of Helsinki and complied with the Guideline of Bioethics and Safety Act by the Ministry of Health and Welfare. The study was approved by the Institutional Review Board of KC Skin Research Center Co., Ltd., in Korea (KC-IRB-016). 21 healthy female subjects who aged 30 ~ 60 years and who have voluntarily signed the Consent form after understanding study object and contents, and who can be observed and traced during the study period. In vivo test was carried out during the period of four weeks with test subjects. A total of twenty female subjects were selected after homogeneity test. The individual instructions over how to use each product are as follows. First, subjects were instructed to use "Biotoc Regen" ampoule, serum and cream individually for four weeks, by applying a proper amount of the product onto their faces and patting the contents to be absorbed, twice a day (morning and night). Second, the subjects were instructed to use "Biotoc Regen" peel individually for four weeks, once a week (evening). The evaluation of adverse events was carried out by the investigator during every visit during the skin condition measurement and analysis activities. Erythema, edema, scaling, itching, stinging, burning sensation, tightness, prickling, and other abnormalities were evaluated. Severities were classified as weak,

moderate, or severe. The test was recorded in the case record by checking whether the test was stopped or omitted. If a subject was no longer able to participate in the examinations even though it was not a visiting day, they would have to fill out the "Attendance Abandonment Agreement" and enclose their signature. To minimize errors in the measurement of the skin conditions of the subjects, the subjects first rinsed their skin with the same cleanser and then stabilized their skin for 30 minutes in a constant temperature and humidity chamber (temperature: $22 \pm 1^\circ\text{C}$, humidity: $45\% \pm 5\%$). Crow's feet, skin moisturization, and TEWL were measured. Measuring instruments used were the most used instruments for skin wrinkles (Antera 3D, Miravex Ltd, Ireland), moisturization (Corneometer, Courage Inc, Germany), and TEWL (Vapometer, Courage Inc, Germany).

Crow's feet were measured by using Antera 3D to photograph crow's feet before using the test product, 2 weeks after using the product, and 4 weeks after using the product. A specific area of the photograph was specified and analyzed and used as an evaluation material for crow's feet. The analyzed wrinkle variable is the overall size, and since the analysis value and the improvement degree of wrinkle are inversely proportional, it means that the lower the analysis value, the better the wrinkle.

Skin moisturization was measured by using Corneometer before using the test product, 2 weeks after using the product, and 4 weeks after using the product. The skin moisture content of the facial cheeks was measured. It was measured 3 times and averaged to evaluate skin moisture content. The unit of skin moisture content is Arbitrary Unit (A.U.), and the measured value is proportional to the skin moisture content, it means that the higher the measured value, the better the skin moisturization.

TEWL was measured in the facial cheeks by using Vapometer before using the test product, 2 weeks after using the product, and 4 weeks after using the product. The value measured once is used as evaluation data, and the unit is g/m²h. Since the measured value and the improvement degree of TEWL are inversely proportional, it means that the lower the measured value, the better the TEWL.

Statistical Analysis

Statistical analysis of the data in the present study was conducted in IBM SPSS Statistics 23.0 for Windows (IBM-Armonk, NY, USA). A paired t-test was used to analyze the significant

changes in the results for various skin characteristics. Differences were accepted as statistically significant at $p < 0.05$

Results and Discussions

Crow's Feet Changes of the Subject

The changes in the crow feet were measured 3 times, including before application, after 2 weeks of use, and after 4 weeks of use. Measurement of facial wrinkles after use of the test products showed a decrease to 15.700 ± 4.201 after 2 weeks and 15.417 ± 3.998 after 4 weeks, from 16.995 ± 4.625 (Figure 1, $p < 0.05$). According to the improvement rate analysis, the degree of wrinkles decreased by 7.199% after 2 weeks, 8.528% after 4 weeks (Table 1). In other words, the analysis of wrinkles after using the test products revealed a statistically significant decrease 2 weeks after using the products and 4 weeks after using the products ($p < 0.05$). Therefore, it was concluded that using the test products facilitated the improvement of wrinkles.

Table 1
Results of Crow's Feet Changes

Time	Average \pm STD (Overall size)	Improvement rate ^a (%)	Probability ^b (p value)
Before use	16.995 ± 4.625	-	-
After 2 weeks of use	15.700 ± 4.201	-7.199	0.000*
After 4 weeks of use	15.471 ± 3.998	-8.258	0.000*

Improvement rate^a (%) = [(after product use – before product use) / before product use] x 100

Probability^b (p value) *: $p < 0.05$ by paired samples T-test




Figure 1. Results of crow's feet changes using Antera 3D

Measurement Results of Skin Moisturization

The changes in the skin moisturization were measured 3 times, before using the test products, 2 weeks after using the products, and 4 weeks after using the products. The results showed that the moisturization increased from 60.383 ± 8.466 to 65.632 ± 8.409 after 2 weeks and to 65.162 ± 8.113 after 4 weeks (Figure 2, $p < 0.05$). The degrees of the improvement in the 2 and 4 weeks were increased by 9.013% after 2 weeks, while increased by 8.287% after 4 weeks (Table 2). According to the results, measurement results of skin moisturization after the test products use, measured value of skin moisture content statistically significantly increased 2 weeks after and 4 weeks after use of products compared to before use of products ($p < 0.05$).

Table 2

Results of Skin Moisturization Measurements

Time	Average \pm STD (A.U.)	Improvement rate ^a (%)	Probability ^b (p value)
Before use	60.383 ± 8.466	-	-
After 2 weeks of use	65.632 ± 8.409	9.013	0.000*
After 4 weeks of use	65.162 ± 8.113	8.287	0.000*

Improvement rate^a (%) = [(after product use – before product use) / before product use] x 100

Probability^b (p value) *: $p < 0.05$ by paired samples T-test




Figure 2. Results of skin moisturization using Corneometer

Measurement Results of TEWL

Changes in TEWL were assessed 3 times, including before using products, after 2 weeks, and after 4 weeks of using products. Using the test products resulted in a decrease of the index from 14.048 ± 2.539 before use, to 13.005 ± 2.472 after 2 weeks of use and to 13.090 ± 2.510 after

four weeks (Figure 3, $p < 0.05$). Analysis of the rate of improvement in TEWL revealed that the TEWL decreased by 7.306% after 2 weeks and 6.559% after 4 weeks (Table 3). Therefore, measurement results of TEWL after the test products use, measured value of TEWL statistically significantly decreased 2 weeks after and 4 weeks after use of products compared to before use of products ($p < 0.05$).

Table 3
Results of TEWL

Time	Average \pm STD (g/m ² h)	Improvement rate ^a (%)	Probability ^b (p value)
Before use	14.048 \pm 2.539	-	-
After 2 weeks of use	13.005 \pm 2.472	-7.306	0.001*
After 4 weeks of use	13.090 \pm 2.510	-6.559	0.005*

Improvement rate^a (%) = [(after product use – before product use) / before product use] x 100

Probability^b (p value) *: $p < 0.05$ by paired samples T-test




Figure 3. Results of TEWL using Vapometer

Evaluation of Skin Adverse Reactions

The presence of skin adverse reactions such as erythema, edema, scaling, itching, stinging, burning, tightness, and pricking was investigated. No specific skin adverse events were observed in all subjects participated in the present study (Table 4).

Table 4
Assessing Skin Adverse Events

Time	Erythema	Edema	Scaling	Itching
After 2 weeks	-	-	-	-

After 4 weeks	-	-	-	-
Time	Stinging	Burning	Tightness	Pricking
After 2 weeks	-	-	-	-
After 4 weeks	-	-	-	-

Step = 1: Weak, 2: Medium, 3: Severe

Conclusions

This study focused on verifying whether the use of a cosmetics containing 1,000ppm Pal-GHK-R4 has a positive effect on facial skin. *In vivo* test was carried out during the period of 4 weeks with test subjects. A total of 21 female subjects were selected after homogeneity test. The individual instructions over how to use each product are as follows. First, subjects were instructed to use Biotoc Regen ampoule, serum, and cream individually for 4 weeks, by applying a proper amount of the products onto their faces and patting the contents to be absorbed, twice a day. Second, the subjects were instructed to use Biotoc Regen peel individually for 4 weeks, once a week. Subjects were asked to wash with their facial skin the same cleanser and after 30 minutes of stabilization in an indoor environment maintained at constant temperature and humidity, 3 kinds of skin testers; Antera 3D for crow's feet, Corneometer for skin moisturization, and Vapometer for TEWL were used to evaluate different facial characteristics.

When the test products were applied, changes of the crow's feet, skin moisturization and TEWL were analyzed. It was observed that crow's feet decreased by 7.199% after 2 weeks of use and 8.258% after 4 weeks of use. The test product had a very significant effect on the improvement of crow's feet. In the case of skin moisturization, it increased by 9.013% 2 weeks after products use and 8.287% 4 weeks after products use. The test product had a very significant positive effect on skin moisturization. Finally, in the case of TEWL, it decreased by 7.306% 2 weeks after using products and 6.559% 4 weeks after using products. The test product also had a very significant positive effect on TEWL. Therefore, the test products containing wrinkle improving peptides, Pal-GHK-R4, are considered to have beneficial effects for 4 weeks of use on improvement of crow's feet, skin moisturization, and TEWL.

Acknowledgment

This study was supported by the Bio & Medical Technology Development Program of the National Research Foundation (NRF) funded of the Ministry of Science & ICT (2017M3A9D8048416).

References

1. Yang, Y. (2010). The Responding Strategies of Musculoskeletal Disease in Aging Society. *Journal of the Ergonomics Society of Korea*, 29(4), 505-511.
2. Oh, J. S. (2017). A Study on the Usage and Purchase Intention of Basic Cosmetics for Silver Generation Women. *J. Cult. Exchange*, 6(3), 77-97.
3. Kang, K. S., Kim, I. D., Kwon, R. H., Heo, Y. Y., Oh, S. H., Kim, M. A., ... & Ha, B. J. (2007). The evaluation of anti-wrinkle effects in oriental herb extract. *Journal of Life Science*, 17(8), 1147-1151.
4. Moh, S. H., Jung, D. H., Kim, H. S., Cho, M. J., Seo, H. H., & Kim, S. J. (2011). Characteristics and applications of bioactive peptides in skin care. *KSBB Journal*, 26(6), 483-490.
5. Yoo, B. J., & Yeon, J. J. (2017). Hwajangpum Wonlyoyong Peptaideu-ui Hyomo Balhyo Saengsan Gisul. *BT News*, 24(2), 52-56.
6. Li, H., Low, Y. S. J., Chong, H. P., Zin, M. T., Lee, C. Y., Li, B., ... & Kang, L. (2015). Microneedle-mediated delivery of copper peptide through skin. *Pharmaceutical research*, 32(8), 2678-2689.
7. Maquart, F. X., Pickart, L., Laurent, M., Gillery, P., Monboisse, J. C., & Borel, J. P. (1988). Stimulation of collagen synthesis in fibroblast cultures by the tripeptide-copper complex glycyl-L-histidyl-L-lysine-Cu²⁺. *FEBS letters*, 238(2), 343-346.
8. Pickart, L. (2008). The human tri-peptide GHK and tissue remodeling. *Journal of Biomaterials Science, Polymer Edition*, 19(8), 969-988.
9. Maquart, F. X., Bellon, G., Chaqour, B., Wegrowski, J., Patt, L. M., Trachy, R. E., ... & Gillery, P. (1993). In vivo stimulation of connective tissue accumulation by the tripeptide-copper complex glycyl-L-histidyl-L-lysine-Cu²⁺ in rat experimental wounds. *The Journal of clinical investigation*, 92(5), 2368-2376.
10. Godet, D., & Marie, P. J. (1995). Effects of the tripeptide glycyl-L-histidyl-L-lysine copper complex on osteoblastic cell spreading, attachment and phenotype. *Cellular and molecular biology (Noisy-le-Grand, France)*, 41(8), 1081-1091.
11. Pickart, L., Freedman, J. H., Loker, W. J., Peisach, J., Perkins, C. M., Stenkamp, R. E., & Weinstein, B. (1980). Growth-modulating plasma tripeptide may function by facilitating copper uptake into cells. *Nature*, 288(5792), 715-717.
12. Lee, H. J., Huang, Y. W., Chiou, S. H., & Aronstam, R. S. (2019). Polyhistidine facilitates direct membrane translocation of cell-penetrating peptides into cells. *Scientific reports*, 9(1), 1-11.

13. Vives, E., Brodin, P., & Lebleu, B. (1997). A truncated HIV-1 Tat protein basic domain rapidly translocates through the plasma membrane and accumulates in the cell nucleus. *Journal of Biological Chemistry*, 272(25), 16010-16017.
14. Wang, K., Zhao, X., Yang, F., Liu, P., & Xing, J. (2019). Percutaneous Delivery Application of Acylated Steric Acid-9-P (arginine) Cell Penetrating Peptides Used as Transdermal Penetration Enhancer++. *Journal of biomedical nanotechnology*, 15(3), 417-430.
15. Yi, D., Guoming, L., Gao, L., & Wei, L. (2007). Interaction of arginine oligomer with model membrane. *Biochemical and biophysical research communications*, 359(4), 1024-1029.
16. Liang, W., Davalian, D., & Torchilin, V. P. (2004). Interaction of a novel peptoid enhancer--arginine oligomer with bovine submaxillary mucin. *Yao xue xue bao= Acta pharmaceutica Sinica*, 39(12), 1011-1017.
17. Mazurowska, L., & Mojski, M. (2008). Biological activities of selected peptides: skin penetration ability of copper complexes with peptides. *Journal of cosmetic science*, 59(1), 59-69.
18. Koch, P. J., De Viragh, P. A., Scharer, E., Bundman, D., Longley, M. A., Bickenbach, J., ... & Roop, D. R. (2000). Lessons from loricrin-deficient mice: compensatory mechanisms maintaining skin barrier function in the absence of a major cornified envelope protein. *The Journal of cell biology*, 151(2), 389-400.