Propolis In Dentistry – A Review

Turkish Online Journal of Qualitative Inquiry (TOJQI) Volume 6, July 2021: 6307- 6316

Research Article

Propolis In Dentistry – A Review

Dr Venkatesh Kodgi ^a, Dr Nithya Annie Thomas ^b, Dr Priya Shetty ^c, Dr Shridhar Shetty ^d, Dr Bekal Kavita Kriplani ^e, Dr Parimala Kumar ^f

^a Chief consultant pediatric dentist, Pearl dental specialty clinic, Muscat, Sultanate of Oman
 ^b Senior Lecturer, Department of Pediatric and Preventive Dentistry, Indira Gandhi Institute of dental sciences, Nellikuzhi P.O., Kothamangalam, Ernakulam P.O., Kerala, India
 ^c Associate Professor, Department of Pediatric and Preventive Dentistry, A.J. Institute of dental sciences, Kuntikana P.O., Mangalore, Karnataka, India
 ^d Professor & HOD, Department of Pediatric and Preventive Dentistry, Yogita dental college and hospital, Khed, Maharashtra, India
 ^e Chief consultant pediatric dentist, Dr. LH Hiranandani hospital, Hillside road, Hiranandani Gardens, IIT Area, Powali, Mumbai, Maharashtra, India
 ^f Associate Professor, Department of Periodontics, A.J. Institute of dental sciences, Kuntikana P.O., Mangalore, Karnataka, India

b*Corresponding author Email- nithyaannie@gmail.com

Abstract

Use of natural products with medicinal properties, derived from plants, animals or microorganisms has found significant popular acceptance in recent years. 'Propolis' is one such natural substance, a resin procured from honeybees collected from plants, buds and exudates has a spectrum of effects like antiviral, antioxidant, antitumor, antibacterial, antifungal, antitumor and immune enhancing properties. Considering oral health, propolis has been utilised in the management of dental caries, pulpitis, aphthous ulcers, plaque, periodontitis, halitosis, stomatitis, dentinal hypersensitivity, oral candidiasis, acute necrotizing ulcerative gingivitis (ANUG), pulp therapy, intra canal medicament and in oral cancer therapy. Pubmed, Google scholar and Medline databases were searched for articles related to propolis and dentistry, which is presented in this review.

Keywords: Propolis, Dentistry, Oral health

1. Introduction

Phytomedicine is increasingly gaining a key role in the dental arena with many plant-derived materials being researched upon. The use of products derived from natural sources such as plants, microorganisms or animals recorded in history dated back to centuries. Since our forefathers chewed on certain herbs for pain relief, natural products were the exclusive source to treat diseases and injuries.

'Propolis', one such natural product, is a resin obtained from bees (*Apis mellifera*) from materials collected from plants, exudates and buds ^{1,2}. The term propolis is derived from the Greek word 'pro' (in front of) and 'polis' (community), interpreting that this product is associated with hive defense^{3,4}. Propolis was customary to Egyptian high priests who had monopolized the proficiency of mummifying corpses ^{1,5}. The initial scientific work with propolis, was published in 1908⁶. The therapeutic use of products procured from honeybees is called Apitherapy ⁷. In the past 30 years, extensive scientific research had been performed to clarify the medicinal properties of this material^{8,9}.

Propolis is a composite mixture of more than 300 naturally occurring constituents identified to date mostly of resins, fatty acids, aromatic oils, pollen, bee wax, vitamins and minerals^{10,11,12}. Its chemical composition includes polysaccharides, phenols, terpenes, aldehydes, tannins, aromatic acids and many more compounds ^{13,14}. Its composition varies with the species of bees, botanical source, geography, and climate, season. In temperate belt, the bud exudates of Populus species along with its hybrids are the primary source of Proplis. This is true for parts of Asia and Europe^{2,15,16}. The most common type of tropical propolis is the Brazilian green propolis^{17,18}.

The biologically activite constituents of propolis are flavonoids, chrysin, caffeic acid, terpenoids, apigenin, pinocembrin, phenolic acids, pinobanksin, pinobanksin 3-acetate, pinobanksin 5-methyl ether ^{19,20}.

Propolis is available in the raw form, tinctures, pills, semisolid preparations, mixtures, emulsions, concentrates, creams, ointments. In dentistry, it is used in toothpastes, mouth rinses, oral sprays and creams. The medicinal properties of this wonder material have rendered it of great scientific interest in recent years.

2. Potential uses of Propolis In Dentistry

Pubmed, Google scholar and Medline databases were searched for articles related to propolis and dentistry, which is presented in this review. The various applications of propolis include:

Caries Prevention: Ikeno et. al. (1991), demonstrated the in vitro anticariogenic potential of propolis in rats against Streptococcus mutans, cricetus and sobrinus ²¹. Steinberg et. al. (1996), assessed the antibacterial effect of propolis on oral bacteria and showed that total salivary bacteria and Streptococcus mutans counts were markedly reduced ²². Similarly, Hayacibara M F et. al. (2005) and Duarte et. al. (2006), concluded that propolis inhibited the 'F'-ATPase activity and decreased acid production, demonstrating cariostatic property. Propolis reduced viability of the isolated fractions of mutans streptococci and inhibition of glucosyltransferases activity in rats ^{23,24}. Koru et. al. (2007), found propolis to be effective against several bacteria implicated in caries ²⁵. Duailibe SA et. al. (2007), showed that propolis mouthrinse exhibited antimicrobial activity against Streptococcus mutans in vivo and may be used as an adjunct in the prevention of dental caries²⁶. The antimicrobial effects of various propolis extracts in rats were studied by Arslan S. et. al. (2010). The ethanol and hexane extracts of propolis showed the highest antimicrobial activity²⁷. Kashi et. al. (2011), revealed that the ethanolic extract showed bacteriostatic and bactericidal activity against all the strains whereas the water extract showed bactericidal activity only against S. mutans in vitro²⁸. Liberio SA et. al. (2011), showed that the Propolis extract with the highest concentration of flavonoids, displayed the most powerful antimicrobial activity on oral pathogens and summarised that it can be made use of in the prevention of caries, candidiasis, and treatment of inflammation²⁹. Krumina G et. al. (2014), suggested that all ethanolic extracts demonstrated antibacterial activity against S. mutans³⁰. Franca et. al. (2014), carried out an in-vitro study with a sustained-release propolis impregnated chitosan varnish and chlorhexidine for biofilm formation prevention and found that, the cariogenic microorganisms of the biofilm showed more favourable results with propolis than chlorhexidine and the bioactive components of Propolis were discharged for more than seven days ³¹. Another study showed definite antimicrobial action of propolis against S. mutans and C. albicans, the probable organisms involved in caries progression³². Tulsani SG et. al. (2015), documented about the anticariogenic action of Propolis gum in comparison to Xylitol gum ³³.

Oral candidiasis: The antifungal efficacy of propolis was demonstrated by Ota *et. al.* (2001) on 80 strains of Candida yeasts ³⁴. The inhibitory effect of 20% ethanol propolis extract (EPE) on the C. albicans multiplication collected from HIV positive persons was tested in vitro³⁵. D'Auria FD *et. al.* (2003), attributed the effects of propolis against C. albicans to be rapid, dose- dependent cytocidal activity and an inhibitory effect on yeast-mycilial conversion³⁶. Patients with oral candidiasis treated with propolis had similar outcome as those treated with nystatin³⁷. Sinha DJ *et. al.* (2015), compared the effect of various herbal extracts and propolis on Candida albicans biofilm, on extracted human mandibular premolars and suggested that, sodium hypochlorite and propolis groups exhibited high antimicrobial efficacy against C. albicans and propolis was better than tea tree oil, turmeric and neem extracts³⁸.

Stomatitis and mouth ulcers: Denture stomatitis is a very common inflammatory lesion on the palatal mucosa of denture wearers. Studies indicate that administration of propolis significantly reduced the outbreaks of recurrent aphthous stomatitis ^{39,40}. Denture stomatitis also responded well to propolis gel in a study and warranted to a complete clinical remission of stomatitis ^{[41].} On the contrary, Da Silva *et. al.* (2008), concluded that a gel containing propolis, can have an adverse effect on the acrylic resin surface which becomes rough favouring microbial adhesion ⁴². It was also found that Brazilian green propolis was beneficial in the management of Candida-associated denture stomatitis ⁴³.

Plaque control: Murray MC et. al. (1997), analysed the effectiveness of a propolis mouthrinse in the plaque

inhibition and observed that chlorhexidine mouthrinse was superior in plaque inhibition ⁴⁴. Giamalia *et. al.* (1999), found a steady increase in micro hardness of human enamel with increasing concentration of propolis from 0.4% - 2%⁴⁵. This was in accordance with other studies^{46,47}. Koo *et. al.* (2002), studied the efficacy of a propolis containing mouthrinse on a 3-day dental plaque accumulation and indicated that, the propolis mouthrinse was efficacious in inhibiting plaque formation⁴⁸. Similar finding was seen in a study by Dodwad V *et. al.* (2011)⁴⁹. Flavonoids in propolis may aid in the inhibition of calcium phosphate formation⁵⁰. Netto *et. al.* (2013), demonstrated that non-alcoholic propolis may have increased efficiency than 0.12% chlorhexidine when used as a mouthrinse because, after 28 days of use, it suppressed the levels of *Lactobacilli and Mutans streptococci* with better patient acceptability⁵¹. Hegde KS *et. al.* (2013), evaluated the antibacterial activity of a propolis extract in children and suggested it as an alternative antimicrobial agent to prevent plaque and caries⁵². Two other studies showed clinical effectiveness of a toothpaste and gel containing 3% ethanolic extract of propolis in patients with increased risk of gingivitis ^{53,54}. Pedrazzi V *et. al.* (2015), studied the antiplaque activity of one of the plant sources of propolis origin⁵⁵. Ercan, *et. al.* (2015) found that the gingival and plaque indices of the propolis chewing gum group were higher than the propolis mouthwash group ⁵⁶.

Dentinal hypersensitivity (DH): This condition is a sharp pain arising due to dentinal tubule stimulation. Mahmoud *et. al.* (1999), in their pioneer study found that Propolis was favourable in controlling dentinal hypersensitivity. Flavonoids in propolis, through chelation of metal ions, give rise to dentin desensitization ⁵⁷. Geiger *et. al.* and Peres S. *et. al.*, stated that propolis inhibits the permeability of dentin by 85%^{58,59}. When compared to CPP-ACPF (Casein Phospho Protein- Amorphous Calcium Phosphate Fluoride) and Sodium Fluoride, Propolis performed better in terms of dentin desensitizing⁶⁰. Compared to hydroxyapatite, sodium fluoride and potassium nitrate Propolis reduced dentin hypersensitivity to the maximum extent as studied over a period of 7 days⁶¹.

Chronic Periodontitis: A study by Cairo de Maral (2006) where in the subjects with periodontitis and gingivitis were instructed to brush daily with propolis, and rinse with a propolis solution and apply in periodontal pockets once in 7 days for a period of five weeks demonstrated drop in gingivitis by 95%⁶². Propolis incorporated into a mucoadhesive hydrophilic gel can be useful in treatment of periodontitis⁶³. In an animal study, the systemic administration of propolis reduced the bone loss due to periodontitis⁶⁴. Propolis extract as a subgingival irrigant was more beneficial than scaling and root planing⁶⁵. Vinod KR *et. al.* (2015) observed excellent mucoadhesion due to addition of propolis, vitamin C and E. Propolis gel combination was stable for eight cycles with no agglomeration tendency, and causes augmentation of the contact time of the medicament with that of the mouth⁶⁶.

Storage media for avulsed teeth: Storage medium is an important factor that prevent damage to the periodontal ligament cells. Martin (2004) showed that propolis can be a substitute to Hanks balanced salt solution in order to maintain the viability of the PDL cells after avulsion⁶⁷. Propolis was recommended as a transport medium by Al-Shaher *et. al.*⁶⁸. Ozan *et. al.* (2007), showed that 10% propolis was an effective storage medium⁶⁹. Based on animal studies, it is proposed that, a combination of systemic antimicrobials and propolis could aid in the inhibition of inflammatory root resorption when used as a root surface treatment media^{70,71}. Several studies have shown that propolis and its combinations performed better than milk, egg white and saliva^{72,73,74}. Propolis also reduces the apoptosis of periodontium cells⁷⁵.

Pulp Therapy/ Dentinal Bridge Formation and Pulp Capping: Propolis has shown promising results in inducing a hard tissue barrier after pulp exposure. Bretz *et.al.* (1998), stated that direct capping with calcium hydroxide-based products and propolis, both materials offer an indistinguishable rate of healing of the pulpal inflammation, reducing the pathogenic microbial count and stimulating dentin bridge formation⁷⁶. Stimulation of various enzyme systems and collagen formation could contribute to hard tissue bridge establishment by Propolis⁷⁷. Flavonoids derived from propolis, when used as a pulp capping agent, aid in partial dentinal bridge formation at four weeks⁷⁸. Brazilian propolis paste combined with calcium hydroxide inhibited the growth of the microbes from in vitro primary root canal cultures. The ethanolic extract of propolis was able to produce larger inhibition zones compared to the non-ethanolic extract of propolis according to Rezende *et. al.* (2006). Propolis was comparable to Dycal and MTA as a pulp-capping agent in relation to hard tissue formation⁸⁰. Propolis causes stimulation of stem cells in the absence of pulpal inflammation, infection or necrosis⁸¹.

Obturating agent in primary teeth: Brazilian Propolis incorporated into calcium hydroxide showed larger growth inhibition zones against primary root canal microbe samples than Calcium hydroxide in vitro⁸². Propolis diffused through dentin is utilised as a vehicle for calcium hydroxide⁸³. Because of its effect on the periodontal cells, propolis can be used as a root canal disinfectant^{68,84}. Effectiveness of glycolic propolis as intracanal medicament against E. faecalis, E. coli and endotoxins in root canals was evaluated by Maekawa LE *et. al.* (2013)⁸⁵. Verma MK *et. al.* (2014), confirmed antimicrobial efficacy of 25% aqueous extract of propolis in the primary teeth root canals⁸⁶.

Irrigant and intra-canal medicament: Propolis was evaluated as an endodontic irrigant because of its antibacterial action. Al-Qathami (2003), indicated that the propolis and sodium hypochlorite have similar has antimicrobial activity⁸⁷. Tandon S (2008), demonstrated that propolis is as effective as calcium hydroxide when used as a pulp capping agent ⁸⁸. Several studies concluded that propolis is effective as an intracanal medicament in effective elimination of E.faecalis⁸⁹⁻⁹². Nara A *et. al.* (2010) found that, when used as an irrigant in permanent teeth, 3% sodium hypochlorite and Brazilian propolis showed equal efficacy against E. faecalis⁹³. Guajardo C *et. al.* (2011) demonstrated that calcium hydroxide was more effective than propolis up to 24 hours from application⁹⁴. Studies have demonstrated that propolis, when used in conjunction with passive ultrasonic irrigation is effective in the removal of E. faecalis biofilm from root canal wall^{95,96}. Ramani N *et. al.* (2012) suggested that the alcoholic extract of propolis is effective intracanal medicament⁹⁷. Pimenta HC *et. al.* (2015) showed that, 40% brown propolis paste and calcium hydroxide pastes were effective as an intracanal medicament, against Enterococcus faecalis⁹⁸. However, Jodhka S. *et. al.* (2015) found that propolis demonstrated antimicrobial activity only after 3 days and 2% chlorhexidine gel was a more effective intracanal medicament⁹⁹.

Oral wound healing: Local application of propolis hydroalcoholic solution had no effect on wound healing but accelerated epithelial repair after tooth extraction¹⁰⁰. Magro-Filho (1994) stated that hydroalcoholic mixture of Propolis helped in epithelial repair but had no effect on the wound healing¹⁰¹. The mechanism to hasten wound healing by propolis is through the activity of caffeic acid phenyl ester (CAPE) that enhances the production of submucosal collagen¹⁰². Propolis decreases inflammation and accelerates epithelialisation and formation of granulation tissue¹⁰³. Studies conducted on rats showed increased amounts of collagen following topical application of propolis on oral ulcers and increased quantity of osteoblasts and rapid remodeling within the palatine suture during bone forming process^{104,105}.

Oral cancer treatment: Animal studies suggest that propolis aid in the chemically induced lingual carcinogenesis in rats and several hydro soluble compounds of propolis, such as CAPE could be useful for the management of tumour growth in pilot studies^{106,107}. CAPE could be used as an agent to prevent oral cancer metastasis¹⁰⁸. CAPE is also an adjuvant for the management of oral squamous cell carcinoma ¹⁰⁹.

3. Safety Issues and Drug Interactions

Phytomedicines although naturally derived, cannot always be regarded as safe. Complete knowledge of the biocompatibility should be present, prior to clinical usage. Allergens have been isolated from propolis. Adverse allergic reactions may manifest as peeling of lips, contact chelitis, perioral eczema, contact stomatitis, oral pain, dyspnea and labial edema¹¹⁰. Presentation may be in the form of redness of skin, rashes, swelling, itching and fever¹¹¹. Allergies related to propolis containing toothpaste have increased from 0.5% to 1.4%¹¹². Routine patch testing should be done in patients before prescribing it. Propolis is said to have interactions with other drugs like metronidazole. Propolis may interact with antifungals, H. pylori agents, anticoagulants, antibiotics, anti-cancer agents, anti-inflammatories, immunosuppressants and antiretrovirals¹¹⁰.

4. Conclusion

Although most of the literature published on propolis is based on animal studies or *in-vitro* studies, extrapolating the result of these studies to clinical practice may not be necessarily replicated in human trials. While using this "api-herbal drug", the clinician should be vigilant due to its hypersensitivity issues shown in a few cases. Evidence from randomized control trials and controlled clinical trials are lacking. Further studies need to be conducted to determine its effects on the oral cavity. In the subsequent years, propolis may find a definitive and promising contribution in dentistry..

References

- [1] Ghisalberti EL. Propolis: a review. Bee world. 1979 Jan 1;60(2):59-84.
- [2] Greenaway W, Scaysbrook T, Whatley FR. The composition and plant origins of propolis: a report of work at Oxford. Bee world. 1990 Jan 1;71(3):107-18.
- [3] Burdock GA. Review of the biological properties and toxicity of bee propolis (propolis). Food and Chemical toxicology. 1998 Apr 6;36(4):347-63.
- [4] Bankova V, De Castro S, Marcucci M. Propolis: recent advances in chemistry and plant origin. Apidologie. 2000;31(1):3-15.

- [5] Z.A. Makashvili, From the history of propolis. In Remarkable hive product: Propolis. Scientific data and suggestions concerning its composition, properties and possible use in therapeutics. APIMONDIA standing commission on beekeeping technology and equipment, Bucharest, 1978
- [6] Helfenberg KD. The analysis of beeswax and propolis. Chemiker Zeitungm. 1908;31:987-98.
- [7] Hellner M, Winter D, von Georgi R, Münstedt K. Apitherapy: usage and experience in german beekeepers. Evidence-Based Complementary and Alternative Medicine. 2008;5(4):475-9.
- [8] Banskota AH, Tezuka Y, Kadota S. Recent progress in pharmacological research of propolis. Phytotherapy Research. 2001 Nov 1;15(7):561-71.
- [9] Noronha VR. Evidencias preliminares da eficácia de gel contendo propolis na prevenção e tratamento de mucosite e candidose bucais em pacientes submetidos a radioterapia em região de cabeça e pescoço (Doctoral dissertation, Thesis. Minas Gerais Federal University, Belo Horizonte. Brazil. Portuguese).
- [10] Kamburoğlu K, Özen T. Analgesic effect of Anatolian propolis in mice. Agri. 2011 Apr;317(23):2.
- [11] Ozan F, Sümer Z, Polat ZA, Er K, Ozan U, Deger O. Effect of mouthrinse containing propolis on oral microorganisms and human gingival fibroblasts. Eur J Dent. 2007 Oct;1(4):195-201.
- [12] Skaba D, Morawiec T, Tanasiewicz M, Mertas A, Bobela E, Szliszka E, Skucha-Nowak M, Dawiec M, Yamamoto R, Ishiai S, Makita Y. Influence of the toothpaste with brazilian ethanol extract propolis on the oral cavity health. Evidence-Based Complementary and Alternative Medicine. 2013 Jun 4;2013.
- [13] Asis M. Propóleos: el oro púrpura de las abejas. Centro de Información y Documentación Agropecuaria. CIDA, La Habana, Cuba. 1989.
- [14] Koo MH, Park YK. Investigation of flavonoid aglycones in propolis collected by two different varieties of bees in the same region. Bioscience, biotechnology, and biochemistry. 1997 Jan 1;61(2):367-9.
- [15] Bankova V, Dyulgerov A, Popov S, Evstatieva L, Kuleva L, Pureb O, Zamjansan Z. Propolis produced in Bulgaria and Mongolia: phenolic compounds and plant origin. Apidologie. 1992 Jan 1;23(1):79-85.
- [16] Johnson KS, Eischen FA, Giannasi DE. Chemical composition of North American bee propolis and biological activity towards larvae of greater wax moth (Lepidoptera: Pyralidae). Journal of Chemical Ecology. 1994 Jul 1;20(7):1783-91.
- [17] Tazawa S, Warashina T, Tadataka NO, MIYASE T. Studies on the constituents of Brazilian propolis. Chemical and Pharmaceutical Bulletin. 1998 Sep 15;46(9):1477-9.
- [18] Trusheva B, Popova M, Bankova V, Simova S, Marcucci MC, Miorin PL, Pasin FD, Tsvetkova I. Bioactive constituents of Brazilian red propolis. Evidence-Based Complementary and Alternative Medicine. 2006;3(2):249-54.
- [19] Huang S, Zhang CP, Wang K, Li GQ, Hu FL. Recent advances in the chemical composition of propolis. Molecules. 2014 Nov 26;19(12):19610-32.
- [20] Marcucci MC. Propolis: chemical composition, biological properties and therapeutic activity. Apidologie. 1995 Mar;26(2):83-99.
- [21] Ikeno K, Ikeno T, Miyazawa C. Effects of propolis on dental caries in rats. Caries Research. 1991 Jul 1;25(5):347-51.
- [22] Steinberg D, Kaine G, Gedalia I. Antibacterial effect of propolis and honey on oral bacteria. American journal of dentistry. 1996 Dec;9(6):236-9.
- [23] 22 D. Steinberg, G. Kaine, I. Gedalia. Antibacterial effect of propolis and honey on oral bacteria. Am. J. Dent., 9 (1996), pp. 236–239
- [24] [23] Hayacibara MF, Koo H, Rosalen PL, Duarte S, Franco EM, Bowen WH, Ikegaki M, Cury JA. In vitro and in vivo effects of isolated fractions of Brazilian propolis on caries development. Journal of Ethnopharmacology. 2005 Oct 3;101(1):110-5.
- [25] Duarte S, Rosalen PL, Hayacibara MF, Cury JA, Bowen WH, Marquis RE, Rehder VL, Sartoratto A, Ikegaki M, 25 Koru O, Toksoy F, Acikel CH, Tunca YM, Baysallar M, Uskudar Guclu A, Akca E, Ozkok Tuylu A, Sorkun K, Tanyuksel M, Salih B. In vitro antimicrobial activity of propolis samples from different geographical origins against certain oral pathogens. Anaerobe 2007 Jun-Aug; 13(3-4):140-5.

- [26] Duailibe SA, Gonçalves AG, Ahid FJ. Effect of a propolis extract on Streptococcus mutans counts in vivo. Journal of Applied Oral Science. 2007 Oct;15(5):420-3.
- [27] Arslan S, Silici S, Percin D, Koç AN, Er Ö. Antimicrobial activity of poplar propolis on mutans streptococci and caries development in rats. Turkish Journal of Biology. 2012 Jan 30;36(1):65-73.
- [28] Jafarzadeh Kashi TS, Kermanshahi RK, Erfan M, Vahid Dastjerdi E, Rezaei Y, Tabatabaei FS. Evaluating the in-vitro antibacterial effect of Iranian propolis on oral microorganisms. Iranian Journal of Pharmaceutical Research. 2011 Jun 19:363-8.
- [29] Liberio SA, Pereira AL, Dutra RP, Reis AS, Araújo MJ, Mattar NS, Silva LA, Ribeiro MN, Nascimento FR, Guerra RN, Monteiro-Neto V. Antimicrobial activity against oral pathogens and immunomodulatory effects and toxicity of geopropolis produced by the stingless bee Melipona fasciculata Smith. BMC Complementary and Alternative Medicine. 2011 Nov 4;11(1):108.
- [30] Krumina G, Ratkevicha L, Nikolajeva V, Babarikina A, Babarykin D. Influence of plant extracts on the growth of oral pathogens Streptococcus mutans and Candida albicans in vitro. Proc Estonian Acad Sci 2015; 64:62–67
- [31] Franca JR, De Luca MP, Ribeiro TG, Castilho RO, Moreira AN, Santos VR, Faraco AA. Propolis-based chitosan varnish: drug delivery, controlled release and antimicrobial activity against oral pathogen bacteria. BMC complementary and alternative medicine. 2014 Dec 12;14(1):478.
- [32] C. Marya, M. Chopra, S. Oberoi, R. Nagpal and C. Dhingra, Jundishapur J Nat Pharm Prod., 10 (2015.
- [33] Tulsani SG, Chikkanarasaiah N, Siddaiah SB, Krishnamurthy NH. The effect of Propolis and Xylitol chewing gums on salivary Streptococcus mutans count: A clinical trial. Indian Journal of Dental Research. 2014 Nov 1;25(6):737.
- [34] Ota C, Unterkircher C, Fantinato V, Shimizu MT. Antifungal activity of propolis on different species of Candida. Mycoses. 2001 Nov 1;44(9-10):375-8.
- [35] Martins RS, Péreira ES, Lima Jr SM, Senna MI, Mesquita RA, Santos VR. Effect of commercial ethanol propolis extract on the in vitro growth of Candida albicans collected from HIV-seropositive and HIVseronegative Brazilian patients with oral candidiasis. Journal of oral science. 2002;44(1):41-8.
- [36] D'auria FD, Tecca M, Scazzocchio F, Renzini V, Strippoli V. Effect of propolis on virulence factors of Candida albicans. Journal of chemotherapy. 2003 Jan 1;15(5):454-60.
- [37] Santos VR, Pimenta FJ, Aguiar MC, Do Carmo MA, Naves MD, Mesquita RA. Oral candidiasis treatment with Brazilian ethanol propolis extract. Phytotherapy Research. 2005 Jul 1;19(7):652-4.
- [38] Sinha DJ, Vasudeva A, Gowhar O, Garg P, Sinha A, Prakash P. Comparison of antimicrobial efficacy of propolis, Azadirachta indica (Neem), Melaleuca alternifolia (Tea tree oil), Curcuma longa (Turmeric) and 5% sodium hypochlorite on Candida albicans biofilm formed on tooth substrate: An in-vitro study. J Pharm Biomed Sci. 2015;5(6):469-74.
- [39] Samet N, Laurent C, Susarla SM, Samet-Rubinsteen N. The effect of bee propolis on recurrent aphthous stomatitis: a pilot study. Clinical oral investigations. 2007 Jun 1;11(2):143-7.
- [40] Preeti L, Magesh KT, Rajkumar K, Karthik R. Recurrent aphthous stomatitis. Journal of Oral and Maxillofacial Pathology. 2011 Sep 1;15(3):252.
- [41] Santos VR, Gomes RT, Mesquita RA, de Moura MD, França EC, Aguiar EG, Naves MD, Abreu JA, Abreu SR. Efficacy of Brazilian propolis gel for the management of denture stomatitis: a pilot study. Phytotherapy Research. 2008 Nov 1;22(11):1544-7.
- [42] Silva WJ, Rached RN, Rosalen PL, Del Bel Cury AA. Effects of nystatin, fluconazole and propolis on poly (methyl methacrylate) resin surface. Brazilian dental journal. 2008;19(3):190-6.
- [43] Capistrano HM, de Assis EM, Leal RM, Alvarez-Leite ME, Brener S, Bastos EM. Brazilian Green própolis compared to miconazole gel in the treatment of Candida-associated denture stomatitis. Evidence-based complementary and alternative medicine. 2013 May 2;2013.
- [44] Murray MC, Worthington HV, Blinkhorn AS. A study to investigate the effect of a propolis-containing mouthrinse on the inhibition of de novo plaque formation. Journal of Clinical Periodontology. 1997 Nov 1;24(11):796-8.

- [45] Giamalia I, Steinberg D, Grobler S, Gedalia I. The effect of propolis exposure on microhardness of human enamel in vitro. Journal of oral rehabilitation. 1999 Dec 1;26(12):941-3.
- [46] Malhotra N, Rao SP, Acharya S, Vasudev B. Comparative in vitro evaluation of efficacy of mouthrinses against Streptococcus mutans, Lactobacilli and Candida albicans. Oral Health and Preventive Dentistry. 2011 Jan 1;9(3):261.
- [47] Shah Rohit A, Kadav Mitali S, Mitra Dipika K, Rodrigues Silvia V, Pathare Pragalbha N. INTERNATIONAL JOURNAL OF RESEARCH IN DENTISTRY COMPARISON OF ANTIPLAQUE EFFICACY OF MOUTHRINSE CONTAINING PROPOLIS WITH 0.2% CHLORHEXIDINE DIGLUCONATE. 2015
- [48] Koo H, Cury JA, Rosalen PL, Ambrosano GA, Ikegaki M, Park YK. Effect of a mouthrinse containing selected propolis on 3-day dental plaque accumulation and polysaccharide formation. Caries Research. 2002 Dec 12;36(6):445-8.
- [49] Dodwad V, Kukreja BJ. Propolis mouthwash: A new beginning. Journal of Indian Society of Periodontology. 2011 Apr 1;15(2):121.
- [50] Hidaka S, Okamoto Y, Ishiyama K, Hashimoto K. Inhibition of the formation of oral calcium phosphate precipitates: the possible effects of certain honeybee products. Journal of periodontal research. 2008 Aug 1;43(4):450-8.
- [51] Netto CA, Marcucci MC, Paulino N, Anido-Anido A, Amore R, de Mendonça S, Neto LB, Bretz WA. Effects of typified propolis on mutans streptococci and lactobacilli: a randomized clinical trial. Brazilian dental science. 2013 Apr 1;16(2):31.
- [52] Hegde KS, Bhat SS, Rao A, Sain S. Effect of propolis on Streptococcus mutans counts: An in vivo study. International journal of clinical pediatric dentistry. 2013 Apr 10;6(1):22-5.
- [53] Tanasiewicz M, Skucha-Nowak M, Dawiec M, Król W, Skaba D, Twardawa H. Influence of hygienic preparations with a 3% content of ethanol extract of brazilian propolis on the state of the oral cavity. Advances in Clinical and Experimental Medicine. 2012 Jan 1;21(1):81-92.
- [54] Morawiec T, Dziedzic A, Niedzielska I, Mertas A, Tanasiewicz M, Skaba D, Kasperski J, Machorowska-Pieniążek A, Kucharzewski M, Szaniawska K, Więckiewicz W. The biological activity of propolis-containing toothpaste on oral health environment in patients who underwent implant-supported prosthodontic rehabilitation. Evidence-based complementary and alternative medicine. 2013 May 14;2013.
- [55] Kurita Varoli F, Sucena Pita M, Sato S, Issa JP, do Nascimento C, Pedrazzi V. Analgesia Evaluation of 2 NSAID Drugs as Adjuvant in Management of Chronic Temporomandibular Disorders. The Scientific World Journal. 2015 Mar 22;2015.
- [56] Ercan N, Erdemir EO, Ozkan SY, Hendek MK. The comparative effect of propolis in two different vehicles; mouthwash and chewing-gum on plaque accumulation and gingival inflammation. European journal of dentistry. 2015 Apr 1;9(2):272.
- [57] Mahmoud AS, Almas K, Dahlan AA. The effect of Propolis on dentinal hypersensitivity and level of satisfaction among patients from a university hospital, Riyadh, Saudi Arabia. Indian J Dent Res 1999;10:130-7.
- [58] Geiger S, Matalon S, Blasbalg J, Tung MS, Eichmiller FC. The clinical effect of amorphous calcium phosphate (ACP) on root surface hypersensitivity. OPERATIVE DENTISTRY-UNIVERSITY OF WASHINGTON-. 2003 Sep 1;28(5):496-500.
- [59] Sales-Peres SH, Carvalho FN, Marsicano JA, Mattos MC, Pereira JC, Forim MR, Silva MF. Effect of propolis gel on the in vitro reduction of dentin permeability. Journal of Applied Oral Science. 2011 Aug;19(4):318-23.
- [60] Madhavan S, Nayak M, Shenoy A, Shetty R, Prasad K. Dentinal hypersensitivity: A comparative clinical evaluation of CPP-ACP F, sodium fluoride, propolis, and placebo. Journal of Conservative Dentistry. 2012 Oct 1;15(4):315.
- [61] Mehta P, Vimala N, Mandke L. An insight into dentin desensitizing agents-In vivo study. Indian Journal of Dental Research. 2013 Sep 1;24(5):571.

- [62] Renata Cairo do Amaral, Rafael Tomaz Gomes, Wellington Márcio Santos Rocha, Sheila Lemos Rago Abreu and Vagner Rodrigues Santos. Periodontitis treatment with brazelian green propolis gel. Pharmacologyonline 3: 336-341 (2006)
- [63] Bruschi ML, Jones DS, Panzeri H, Gremião MP, De Freitas O, Lara EH. Semisolid systems containing propolis for the treatment of periodontal disease: in vitro release kinetics, syringeability, rheological, textural, and mucoadhesive properties. Journal of pharmaceutical sciences. 2007 Aug 1;96(8):2074-89
- [64] Toker H, Ozan F, Ozer H, Ozdemir H, Eren K, Yeler H. A morphometric and histopathologic evaluation of the effects of propolis on alveolar bone loss in experimental periodontitis in rats. Journal of periodontology. 2008 Jun;79(6):1089-94.
- [65] A.O. Coutino, S. Sanikop, KSDJ, 28, 31-35 (2009)
- [66] Vinod KR, Jasmin F, Park JW, Akalin FA, Choi DJ. ROLE OF PROPOLIS IN AUGMENTING THE BUCCAL MUCOADHESION-AN EXPERIMENT BASED REPORT.
- [67] Martin MP, Pileggi R. A quantitative analysis of Propolis: a promising new storage media following avulsion. Dental traumatology. 2004 Apr 1;20(2):85-9.
- [68] Al-Shaher A, Wallace J, Agarwal S, Bretz W, Baugh D. Effect of propolis on human fibroblasts from the pulp and periodontal ligament. Journal of endodontics. 2004 May 31;30(5):359-61.
- [69] Özan F, Polat ZA, Er K, Özan Ü, Değer O. Effect of propolis on survival of periodontal ligament cells: new storage media for avulsed teeth. Journal of Endodontics. 2007 May 31;33(5):570-3.
- [70] Gulinelli JL, Panzarini SR, Fattah CM, Poi WR, Sonoda CK, Negri MR, Saito CT. Effect of root surface treatment with propolis and fluoride in delayed tooth replantation in rats. Dental Traumatology. 2008 Dec 1;24(6):651-7.
- [71] Mori GG, Nunes DC, Castilho LR, Moraes IG, Poi WR. Propolis as storage media for avulsed teeth: microscopic and morphometric analysis in rats. Dental Traumatology. 2010 Feb 1;26(1):80-5.
- [72] Ahangari Z, Alborzi S, Yadegari Z, Dehghani F, Ahangari L, Naseri M. The effect of propolis as a biological Storage media on periodontal ligament cell survival in an avulsed tooth: an in vitro study. Cell J. 2013; 15 (3): 244-249. Introduction Avulsion is defined as the total displacement of a tooth from the alveolar socket. Clinical surveys indi-cate that avulsion occurs in 1to. 2013 Jan 1;16.
- [73] Saxena P, Pant VA, Wadhwani KK, Kashyap MP, Gupta SK, Pant AB. Potential of the propolis as storage medium to preserve the viability of cultured human periodontal ligament cells: an in vitro study. Dental Traumatology. 2011 Apr 1;27(2):102-8.
- [74] Khademi AA, Saei S, Mohajeri MR, Mirkheshti N, Ghassami F. Torabi nia N, Alavi SA. A new storage medium for an avulsed tooth. J Contemp Dent Pract. 2008;9(6):25-32.
- [75] Gjertsen AW, Stothz KA, Neiva KG, Pileggi R. Effect of propolis on proliferation and apoptosis of periodontal ligament fibroblasts. Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology. 2011 Dec 31;112(6):843-8.
- [76] Bretz WA, Chiego DJ, Marcucci MC, Cunha I, Custódio A, Schneider LG. Preliminary report on the effects of propolis on wound healing in the dental pulp. Zeitschrift f
 ür Naturforschung C. 1998 Dec 1;53(11-12):1045-8.
- [77] Park YK, Alencar SM, Aguiar CL. Botanical origin and chemical composition of Brazilian propolis. Journal of Agricultural and Food Chemistry. 2002 Apr 24;50(9):2502-6.
- [78] Sabir A, Tabbu CR, Agustiono P, Sosroseno W. Histological analysis of rat dental pulp tissue capped with propolis. Journal of oral science. 2005;47(3):135-8.
- [79] Rezende GP, Costa LR, Pimenta FC, Baroni DA. In vitro antimicrobial activity of endodontic pastes with propolis extracts and calcium hydroxide: a preliminary study. Brazilian dental journal. 2008;19(4):301-5.
- [80] Parolia A, Kundabala M, Rao NN, Acharya SR, Agrawal P, Mohan M, Thomas M. A comparative histological analysis of human pulp following direct pulp capping with Propolis, mineral trioxide aggregate and Dycal. Australian dental journal. 2010 Mar 1;55(1):59-64.
- [81] Ahangari Z, Naseri M, Jalili M, Mansouri Y, Mashhadiabbas F. Torkaman Pharm A. Effect of propolis on dentin regeneration and the potential role of dental pulp stem cell in Guinea Pigs. Cell J. 2012; 13

(4): 2230228. Cell Journal (Yakhteh). 2012;13(4):223.

- [82] Panzeri H, Lara EH, Pedrazzi V, Gabarra FR, Ito IY. An experimental dentifrice containing propolis: Physical, microbiological and clinical evaluations. InJOURNAL OF DENTAL RESEARCH 1998 Jan 1 (Vol. 77, pp. 975-975). 1619 DUKE ST, ALEXANDRIA, VA 22314 USA: AMER ASSOC DENTAL RESEARCH.
- [83] Montero JC, Mori GG. Assessment of ion diffusion from a calcium hydroxide-propolis paste through dentin. Brazilian oral research. 2012 Aug;26(4):318-22.
- [84] Ramos IF, Biz MT, Paulino N, Scremin A, Della Bona Á, Barletta FB, Figueiredo JA. Histopathological analysis of corticosteroid-antibiotic preparation and propolis paste formulation as intracanal medication after pulpectomy: an in vivo study. Journal of applied oral science. 2012 Feb;20(1):50-6.
- [85] Maekawa LE, Valera MC, Oliveira LD, Carvalho CA, Camargo CH, Jorge AO. Effect of Zingiber officinale and propolis on microorganisms and endotoxins in root canals. Journal of Applied Oral Science. 2013 Feb;21(1):25-31.
- [86] Verma MK, Pandey RK, Khanna R, Agarwal J. The antimicrobial effectiveness of 25% propolis extract in root canal irrigation of primary teeth. Journal of Indian Society of Pedodontics and Preventive Dentistry. 2014 Apr 1;32(2):120.
- [87] Qathami HA, Al-Madi E. Comparison of sodium hypochlorite, propolis and saline as root canal irrigants: A pilot study. Saudi Dental Journal. 2003 Jan 23;15(2):100-3.
- [88] Tandon S. Textbook of pedodontics. Paras Medical Publisher; 2009.
- [89] Kandaswamy D, Venkateshbabu N, Gogulnath D, Kindo AJ. Dentinal tubule disinfection with 2% chlorhexidine gel, propolis, morinda citrifolia juice, 2% povidone iodine, and calcium hydroxide. International endodontic journal. 2010 May 1;43(5):419-23.
- [90] Awawdeh L, AL-Beitawi M, Hammad M. Effectiveness of propolis and calcium hydroxide as a shortterm intracanal medicament against Enterococcus faecalis: A laboratory study. Australian Endodontic Journal. 2009 Aug 1;35(2):52-8.
- [91] Cogulu D, Uzel A, Oncag O, Eronat C. PCR-based identification of selected pathogens associated with endodontic infections in deciduous and permanent teeth. Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology. 2008 Sep 30;106(3):443-9.
- [92] Mattigatti S, Ratnakar P, Moturi S, Varma S, Rairam S. Antimicrobial effect of conventional root canal medicaments vs propolis against Enterococcus faecalis, Staphylococcus aureus and Candida albicans. J Contemp Dent Pract. 2012 May 1;13(3):305-9.
- [93] Nara A, Dhanu PC, Latha Anandakrishna D. Comparative Evaluation of Antimicrobial Efficacy of MTAD, 3% NaOCI and Propolis Against E Faecalis. International Journal of Clinical Pediatric Dentistry. 2010 Jan;3(1):21.
- [94] Cuevas-Guajardo S, Arzate-Sosa G, Flores-Chavez RI, Fabela-Gonzalez LV, Mendieta-Zeron H. Antimicrobial activity with mixture of calcium hydroxide and propolis. International Journal of Pharma and Bio Sciences. 2011;2(4):203-10.
- [95] Bhardwaj A, Velmurugan N, Ballal S. Efficacy of passive ultrasonic irrigation with natural irrigants (Morinda citrifolia juice, Aloe Vera and Propolis) in comparison with 1% sodium hypochlorite for removal of E. faecalis biofilm: An in vitro study. Indian Journal of Dental Research. 2013 Jan 1;24(1):35.
- [96] Bolla N, Kavuri SR, Tanniru HI, Vemuri S, Shenoy A. Comparative evaluation of antimicrobial efficacy of odontopaste, chlorhexidine and propolis as root canal medicaments against enterococcus faecalis and candida albicans. Journal of International Dental and Medical Research. 2012 Apr 1;5(1):14-25.
- [97] Ramani N, Mathew S. Comparative evaluation of antimicrobial efficacy of chlorhexidine digluconate and propolis when used as an intracanal medicament: ex vivo study. Journal of International Oral Health. 2012 May 1;4(2):17.
- [98] Pimenta HC, VIOLANTE IM, MUSIS CR, Borges AH, ARANHA AM. In vitro effectiveness of Brazilian brown propolis against Enterococcus faecalis. Brazilian oral research. 2015;29(1):1-6.

- [99] S. Jodhka, H.S. Bedi, N. Singh, A.M. Thomas, P Ahluwalia, 7, 6-10 (2015)
- [100] FILHO OM, CARVALHO AC. Application of propolis to dental sockets and skin wounds. The Journal of Nihon University School of Dentistry. 1990;32(1):4-13.
- [101] Magro-Filho O, de Carvalho AC. Topical effect of propolis in the repair of sulcoplasties by the modified kazanjian technique. The Journal of Nihon University School of Dentistry. 1994;36(2):102-11.
- [102] Ocakci A, Kanter M, Cabuk M, Buyukbas S. Role of caffeic acid phenethyl ester, an active component of propolis, against NAOH-induced esophageal burns in rats. International journal of pediatric otorhinolaryngology. 2006 Oct 31;70(10):1731-9.
- [103] Lopes-Rocha R, Miranda JL, Lima NL, Ferreira FO, Santos AS, Verli FD, Marinho SA. Effect of topical propolis and dexamethasone on the healing of oral surgical wounds: original research. Wound Healing Southern Africa. 2012 Jan 1;5(1):25-30.
- [104] Moraes LT, Trevilatto PC, Grégio AM, Machado MA, Lima AA. Quantitative analysis of mature and immature collagens during oral wound healing in rats treated by Brazilian propolis. Journal of International Dental and Medical Research. 2011 Dec 1;4(3):106-10.
- [105] Altan BA, Kara IM, Nalcaci R, Ozan F, Erdogan SM, Ozkut MM, Inan S. Systemic propolis stimulates new bone formation at the expanded suture: a histomorphometric study. The Angle Orthodontist. 2012 Aug 20;83(2):286-91.
- [106] Orsolic N, Sver L, Terzic S, Tadic Z, Basic I. Inhibitory effect of water-soluble derivative of propolis and its polyphenolic compounds on tumor growth and metastasizing ability: a possible mode of antitumor action. Nutrition and cancer. 2003 Nov 1;47(2):156-63.
- [107] Cavalcante DR, Oliveira PS, Góis SM, Soares AF, Cardoso JC, Padilha FF, Albuquerque Júnior RL. Effect of green propolis on oral epithelial dysplasia in rats. Brazilian journal of otorhinolaryngology. 2011 Jun;77(3):278-84.
- [108] Peng CY, Yang HW, Chu YH, Chang YC, Hsieh MJ, Chou MY, Yeh KT, Lin YM, Yang SF, Lin CW. Caffeic Acid phenethyl ester inhibits oral cancer cell metastasis by regulating matrix metalloproteinase-2 and the mitogen-activated protein kinase pathway. Evidence-Based Complementary and Alternative Medicine. 2012 Dec 18;2012.
- [109] Kuo YY, Lin HP, Huo C, Su LC, Yang J, Hsiao PH, Chiang HC, Chung CJ, Wang HD, Chang JY, Chen YW. Caffeic acid phenethyl ester suppresses proliferation and survival of TW2. 6 human oral cancer cells via inhibition of akt signaling. International journal of molecular sciences. 2013 Apr 24;14(5):8801-17.
- [110] Basavaiah ND, Suryakanth DB. Propolis and allergic reactions. Journal of pharmacy & bioallied sciences. 2012 Oct;4(4):345.
- [111] Wander P. Taking the sting out of dentistry. Dental practice. 1995;25(1995):3-4.
- [112] Zirwas MJ, Otto S. Toothpaste allergy diagnosis and management. The Journal of clinical and aesthetic dermatology. 2010 May;3(5):42.